



ROTHAMSTED  
RESEARCH

# Inventory of Ammonia Emissions from UK Agriculture 2022

---

Defra Contract C21258  
Inventory Submission Report  
May 2024

A.M. CARSWELL<sup>1</sup>, S.L. GILHESPY<sup>1</sup>, L.M. CARDENAS<sup>1</sup> & S.G. ANTHONY<sup>2</sup>

WITH CONTRIBUTIONS FROM:

**<sup>1</sup>ROTHAMSTED RESEARCH**

A. MILNE, R. SHARP

**<sup>2</sup>ADAS**

H. GRIFFITHS, D. SKIRVIN

**RICARDO**

G. THISTLETHWAITE, K. AVIS, L. PHELPS

**UKCEH**

U. DRAGOSITS, E. CARNELL

**CRANFIELD UNIVERSITY**

D. SANDARS

# CONTENTS

---

<b>Glossary of abbreviations.....</b>	<b>3</b>
<b>Inventory of Ammonia Emissions from UK Agriculture – 2022.....</b>	<b>4</b>
Summary.....	4
Estimate of ammonia emission from UK agriculture for 2022.....	5
Major changes between 2021 and 2022 .....	6
Emission Trends: 1990 - 2022 .....	8
Uncertainties .....	10
<b>Appendix 1: Ammonia Emission Factors for UK Agriculture.....</b>	<b>11</b>
Introduction .....	11
A1.1 Livestock housing .....	11
A1.2 Hard standings (unroofed outdoor concrete yards) .....	13
A1.3 Manure storage.....	13
A1.4 Manure application .....	14
A1.5 Grazing and outdoor livestock.....	16
A1.6 Nitrogen fertiliser applications .....	17
A1.7 Digestate applications to land .....	17
Annex 1: Sources of underlying data for the UK ammonia emission factors....	19
References .....	26
<b>Appendix 2: Reduction efficiencies for ammonia mitigation methods applicable to the UK ammonia emission inventory .....</b>	<b>33</b>
Introduction .....	33
Emission reduction methods.....	33

## GLOSSARY OF ABBREVIATIONS

AFBI	Agri-Food and Biosciences Institute
AHDB	Agriculture and Horticulture Development Board
AS	Ammonium sulphate
BSFP	British Survey of Fertiliser Practice
CAFRE	College of Agriculture, Food and Rural Enterprise
CI	Confidence interval
DA	Devolved Administration
DAERA	Department of Agriculture, Environment and Rural Affairs (Northern Ireland)
DAP	Di-ammonium phosphate
DEFRA	Department for Environment, Food and Rural Affairs (UK)
EF	Emission factor
FYM	Farmyard manure
GHG	Greenhouse Gas
LESSE	Low emission slurry spreading equipment
N	Nitrogen
n	Number of observations
NARSES	National Ammonia Reduction Strategy Evaluation System
NH <sub>3</sub>	Ammonia
NIGTA	Northern Ireland Grain Trade Association
SE	Standard error
TAN	Total ammoniacal nitrogen
UAN	Urea ammonium nitrate
UNECE	United Nations Economic Commission for Europe

# INVENTORY OF AMMONIA EMISSIONS FROM UK AGRICULTURE - 2022

## SUMMARY

The combined UK Agriculture Greenhouse Gas (GHG) and Ammonia emission model was used to compile the 1990-2022 ammonia (NH<sub>3</sub>) emission inventory for UK agriculture, ensuring consistency of approach in terms of nitrogen (N) flows and transformations for both the NH<sub>3</sub> and GHG emission estimates. Year-specific livestock numbers and crop areas were included for 2022, together with information on fertiliser N use for 2022 and any changes in farm management practices where data were available. The estimate of NH<sub>3</sub> emissions from UK agriculture for 2022 was 226.6 kt NH<sub>3</sub>, representing a decrease of 3.97 kt from the previously reported estimate (2023 submission) for 2021. In 2022 agriculture contributed 87% to the total UK NH<sub>3</sub> emissions (Elliot et al., 2024). Updates in the use of N fertiliser activity data, cattle N excretion and manure management activity data resulted in an increase of 0.9 kt in the total estimate for 2021 between the two reporting years. In 2022 NH<sub>3</sub> emissions from agriculture have decreased by 18.9% since 1990 and by 4.4% since 2005.

**Table 1.** Estimate of ammonia emissions from UK agriculture for 2022 with livestock emissions reported either by livestock category (a; which includes emissions from manure-based digestates) or manure management category (b) together with other non-livestock sources (c). Note that table parts a and b have different totals due to the inclusion of manure-based digestate in a.

Source	kt NH <sub>3</sub> <sup>*</sup>	% of total
<b>a. Livestock emissions by livestock category</b>		76.9
Cattle	115.1	50.8
<i>Dairy cows</i>	57.1	25.2
<i>Other cattle<sup>#</sup></i>	57.3	25.3
Sheep	11.9	5.2
Pigs	16.0	7.1
Poultry	30.1	13.3
Minor livestock <sup>†</sup>	1.2	0.5
<b>b. Manure emissions by management category</b>		75.5
Grazing/outdoors	18.8	8.3
Housing	59.5	26.3
Hard standings	14.1	6.2
Manure storage	20.3	8.9
Manure application	58.4	25.8
<b>c. Other sources</b>		24.5
Fertiliser application	34.4	15.2
<i>Urea and UAN fertiliser</i>	22.4	9.9
<i>Other nitrogen fertiliser<sup>§</sup></i>	12.0	5.3
Sewage sludge application	4.7	2.1
Digestate application <sup>  </sup>	16.5	7.3
<i>Non-manure digestate</i>	13.1	5.8
<i>Manure digestate</i>	3.4	1.5
<b>TOTAL</b>	<b>226.6</b>	

<sup>#</sup>Other cattle refers to all beef cattle and non-lactating dairy cattle

---

<sup>†</sup> Horses, goats and deer on agricultural holdings

<sup>\*</sup> Totals may differ from sum of components due to rounding

<sup>§</sup> Other nitrogen fertilisers include ammonium nitrate, calcium ammonium nitrate, ammonium sulphate, diammonium phosphate and other nitrogen fertilisers (including compound blends)

<sup>‡</sup> Manure-based digestate is also included within section a (livestock category totals)

## ESTIMATE OF AMMONIA EMISSION FROM UK AGRICULTURE FOR 2022

The 1990 – 2022 inventory estimates were as made in previous submissions, using the combined GHG and NH<sub>3</sub> emission model for UK agriculture. This model uses the same underlying approach as in the previously used national-scale NARSES model (Webb and Misselbrook, 2004), but incorporates a much higher level of spatial (10 km grid cells), temporal (monthly) and sectoral (greater disaggregation of dairy, beef, sheep, grassland and cropping sectors) resolution for the bottom-up calculations. As part of the model development and improvement, revisions were made to some parameters in the N-flow calculations compared with the NARSES model to ensure consistency between the estimates of NH<sub>3</sub> and GHG emissions. Further details of the model and parameterisation are given in the UK Informative Inventory Report (IIR) and National Inventory Report (NIR) for the 2024 submission.

*Key areas of revision in the 2022 inventory were:*

- Inclusion of 2022 livestock numbers, crop areas and fertiliser N use
- Nitrogen fertiliser activity data was updated with small revisions made to the reporting of estimates of nitrogen fertiliser type used in Northern Ireland, from calendar year to agricultural year; provisional 2021 fertiliser data from the previous submission for Northern Ireland was replaced with actual data; centred three-year rolling averages were applied to the fertiliser rates applied to grass on dairy farm types in Scotland which affects 2020 and 2021
- Minor changes to cattle N excretion associated with updated milk yield and slaughter weight data
- Manure management activity data was updated with changes made to the amount of time spent outside for outdoor poultry, from 20/80 to 10/90 outdoor/indoor; housing emission factor (EF) was updated for weaners and finishers on slatted systems; firm and funded policy from Northern Ireland was implemented for the covering of stores (from 2020) and for increased use of low emission slurry spreading equipment (LESSE); the amount of manure being processed by anaerobic digestion was updated across the time series

Derivations of EFs and reduction efficiencies assumed for mitigation practices are detailed in Appendices 1 and 2.

The estimate of emission from UK agriculture for 2022 was 226.6 kt NH<sub>3</sub>. Cattle represent the largest livestock NH<sub>3</sub> source, with 69.2 kt from housing and storage, 0.1 kt from digestate storage, 36.1 kt from manure applications to land, 1.4 kt from digestate applications to land and 8.4 kt from grazing returns (Table 1a.). Housing and manure application to land are the major NH<sub>3</sub> sources in terms of manure management (Table 1b). A breakdown of the estimate

is given in Table 2, together with a comparison with the previously submitted 2021 inventory estimate.

## **MAJOR CHANGES BETWEEN 2021 AND 2022**

### *1. 2022 livestock numbers*

Headline changes from 2021 were:

Cattle – slight changes in cattle numbers, decrease of 0.4% for dairy cows and increase of 0.5% for other cattle

Pigs – a 3.0% decrease in pig numbers

Sheep – a 0.5% increase in sheep numbers

Poultry – a 2.9% decrease in total poultry numbers, 3.6% decrease in broilers and a 3.5% increase in layers

### *2. Fertiliser N use*

Total fertiliser N use decreased by 16.9% between 2021 and 2022 (Figure 2), with fertiliser applications to tillage decreasing by 8% and to grassland by 30%. The reduction in fertiliser use for 2022 was a result of increased fertiliser prices associated with higher energy prices. In recent years fertiliser use on cropland has varied due to weather conditions, with wet ground conditions disrupting the sowing of winter cereal crops and delaying sowing until the following spring, with spring sown crops requiring less fertiliser. This was observed in 2020, with a 14% decline in total fertiliser N use relative to 2019 and for the years 1992/1993, 2000/2001 and 2013/2014. Typically, fertiliser N use increases again in the following year. The amount of fertiliser N applied as urea and urea ammonium nitrate (UAN) represented 12.6 and 14.7%, respectively, of total fertiliser N use in 2022.

Use of urea-based fertilisers, which are associated with much higher  $\text{NH}_3$  EF than other N fertilisers, has increased as a proportion of total fertiliser N use. In 2021 urea-based fertilisers (including urea and UAN) accounted for 23% of total fertiliser N use and this increased to 27% in 2022 (Figure 2), which is the highest recorded use of urea-based fertilisers in the presented timeseries. Additionally, in 2022 the British Survey of Fertiliser Practice was updated to include foliar urea, which represents a new source of ammonia emissions associated with N fertiliser use. The inclusion of urease inhibitors with urea-based fertilisers reduces  $\text{NH}_3$  emissions, by 70% for urea and 44% for UAN. In 2022 UK uptake of urease inhibitors with urea was 7.5 and 15.3% for arable and grassland crops, respectively, and for UAN was 5.7 and 11.0% for arable and grassland crops, respectively.

In addition to the activity data changes reported for fertiliser N use, some revisions were made to the reporting of estimates of N fertiliser used in Northern Ireland, which changed from a calendar year basis to an agricultural year basis and provisional data used for 2021 in the previous submission was replaced with actual data for this submission. In addition, a centred three-year rolling average was applied to the grassland fertiliser rates for dairy farm types in Scotland, which affected the years 2020 and 2021.

### *3. Change to cattle N excretion rates*

For the 2024 inventory submission (1990 to 2022 time series), small changes to cattle N excretion rates were associated with changes to milk yield (affecting the years 2018 to 2021

for Scotland and Northern Ireland and the years 2019 to 2021 for England and Wales) and the slaughterweight timeseries (from 2008 onwards).

#### 4. Revision to manure management practices

The amount of time that outdoor poultry spend outside was reduced from 20% to 10% based on the outcomes of a review commissioned by the Environment Agency in 2023. Updates were also made to the EF for weaners and finishers on slatted housing systems. Other revisions included the implementation of “firm and funded” policy regarding the covering of stores and increased use of LESSE in Northern Ireland.

**Table 2.** Estimate of ammonia emissions (kt NH<sub>3</sub>) from UK agriculture, 2022\*

Source	2021 as per 2023 submission	2021 as per 2024 submission	Reasons for change between submissions	2022	Reasons for change from 2021
<b>Cattle</b>					
Grazing	8.4	<b>8.4</b>	Slight change in N excretion linked to milk yield changes and slaughter weight updates. Manure storage mitigation introduced for Northern Ireland and amount of slurry spread using LESSE adjusted for all countries. Updated amount of manure diverted to anaerobic digestion.	<b>8.4</b>	Increase in overall cattle numbers. Amount of slurry spread using LESSE increased slightly for all countries
Landspreading	38.3	<b>38.0</b>		<b>37.5</b>	
Housing	42.2	<b>42.2</b>		<b>42.0</b>	
Hard standings	14.1	<b>14.2</b>		<b>14.1</b>	
Storage	13.3	<b>13.3</b>		<b>13.2</b>	
<b>Total Cattle</b>	116.3	<b>116.1</b>		<b>115.1</b>	
<b>Sheep</b>					
Grazing	8.5	<b>8.5</b>	Very small changes in amount of nitrogen excreted associated with changes in amounts of nitrogen fertiliser applied to grass.	<b>8.5</b>	Small increase in sheep numbers
Landspreading	1.3	<b>1.3</b>		<b>1.3</b>	
Housing	1.3	<b>1.3</b>		<b>1.3</b>	
Storage	0.9	<b>0.9</b>		<b>0.9</b>	
<b>Total Sheep</b>	11.9	<b>11.9</b>		<b>11.9</b>	
<b>Minor livestock<sup>†</sup></b>	1.3	<b>1.3</b>		<b>1.2</b>	
<b>Pigs</b>					
Outdoor	1.1	<b>1.1</b>	Changes to amount of nitrogen entering spreading for weaners and EF for weaners and finishers on slatted systems. Manure storage mitigation introduced for Northern Ireland and amount of slurry spread using LESSE adjusted for all countries. Updated amount of manure diverted to anaerobic digestion.	<b>1.0</b>	Decrease in pig numbers
Landspreading	4.3	<b>4.2</b>		<b>4.0</b>	
Housing	8.3	<b>8.2</b>		<b>7.9</b>	
Storage	3.3	<b>3.3</b>		<b>3.2</b>	
<b>Total Pigs</b>	16.9	<b>16.7</b>		<b>16.0</b>	
<b>Poultry</b>					
Outdoor	1.0	<b>0.5</b>	Ratio of outdoor/housed period changed to 10/90 and was previously 20/80. Updated amount of manure diverted to incineration and anaerobic digestion	<b>0.5</b>	Decrease in poultry numbers
Landspreading	18.4	<b>18.8</b>		<b>18.8</b>	
Housing	7.9	<b>8.1</b>		<b>7.9</b>	
Storage	2.8	<b>2.8</b>		<b>2.8</b>	
<b>Total Poultry</b>	30.0	<b>30.2</b>		<b>30.1</b>	
<b>Fertiliser</b>	36.2	<b>37.6</b>	Small revisions made to amounts of N fertiliser applied in Northern Ireland and 2021 estimate updated. Calendar year estimates of N fertiliser use replaced with agricultural year estimates. Dairy farm estimates for Scotland replaced with 3-year average value, which also affected Wales.	<b>34.4</b>	Large decline in N fertiliser application
<b>Sewage sludge</b>	4.7	<b>4.7</b>		<b>4.7</b>	

<b>Non-manure digestate</b>	13.2	<b>12.9</b>	Quantity of digestate updated.	<b>13.1</b>	Increase in the quantity of digestate
<b>TOTAL</b>	230.5	<b>231.5</b>		<b>226.6</b>	

\*Totals may differ from sum of components due to rounding

†Including horses on agricultural holdings, goats and deer

### EMISSION TRENDS: 1990 TO 2022

Retrospective calculations based on the most recent inventory methodology were made for the years 1990 to 2022 (Table 3). There has been a steady decline in emissions from UK agriculture over the period 1990 to 2010, largely due to declining livestock numbers (Figure 1) and N fertiliser use (Figure 2), but also from increases in production efficiency. However, this decline has levelled off in recent years, with increased amounts of non-manure digestate applied to soils since 2005 negating some of the ammonia emission reductions seen for other sources (Table 3). Emissions have declined by 18.9% since 1990, and by 4.4% since 2005, due to a combination of the trend in livestock numbers, fertiliser N use and some uptake of NH<sub>3</sub> abatement techniques. The reduction in NH<sub>3</sub> emissions associated with reduced fertiliser N use in 2022 is not expected to continue if the cost of energy and, consequently, fertiliser decline.


**Table 3.** Estimates of ammonia emission from UK agriculture 1990 – 2022

Source	1990	2000	2005	2010	2015	2020	2021	2022
<b>Total</b>	<b>279.3</b>	<b>248.9</b>	<b>237.1</b>	<b>224.2</b>	<b>237.0</b>	<b>226.7</b>	<b>231.5</b>	<b>226.6</b>
Cattle	112.4	113.2	117.1	114.8	117.7	116.5	116.1	115.1
Sheep	14.8	14.8	12.5	10.6	11.8	11.8	11.9	11.9
Pigs	39.7	29.6	21.2	17.5	16.1	15.9	16.7	16.0
Poultry	51.5	50.3	42.9	33.0	28.4	29.3	30.2	30.1
Minor livestock	1.1	1.5	1.8	1.6	1.5	1.3	1.3	1.2
Fertiliser	51.6	37.7	37.5	41.2	47.8	34.8	37.6	34.4
Sewage sludge	1.5	1.8	3.7	3.9	4.3	4.7	4.7	4.7
Non-manure digestate <sup>1</sup>	0.0	0.0	0.5	1.6	9.4	12.5	12.9	13.1
Field burning	6.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0


<sup>1</sup>Ammonia emission associated with manure-based digestate is included within relevant livestock category



**Figure 1.** Trends in livestock numbers 1990 – 2022, relative to a reference value of 100 in 1990.



**Figure 2.** Changes in fertiliser N use 1990 – 2022.



## UNCERTAINTIES

An estimate of the uncertainties in the emission inventory estimate was conducted using Monte Carlo simulation, in which a probability distribution function was provided for each of the model inputs (activity or EF data), based on the distribution of raw data or, where no or only single estimates exist, on expert assumptions. The 95% confidence interval for the total inventory estimate was estimated to be approximately  $\pm 16.7\%$  (i.e.  $\pm 37.8$  kt  $\text{NH}_3$  for the 2022 estimate).

# APPENDIX 1: AMMONIA EMISSION FACTORS FOR UK AGRICULTURE

## INTRODUCTION

This report described the emission factors (EFs) and where appropriate standard errors (SE) for ammonia (NH<sub>3</sub>) emissions from agricultural sources that are used in the improved greenhouse gas (GHG) emission inventory for UK agriculture, developed under the UK government-funded Defra project AC0114. The improved GHG inventory for UK agriculture uses a nitrogen (N) mass flow approach in calculating emissions from livestock manure management with the initial N input as excretion by livestock and subsequent losses and transformations (between organic and total ammoniacal N, TAN) being modelled at each management stage i.e. livestock housing, manure storage/treatment and manure application to land. Ammonia EFs are expressed as a percentage of the TAN content of the manure N pool at each management stage. In addition, EFs are described for emissions from grazing returns (expressed as a percentage of TAN, which is generally equated with the urine fraction of the excreta) and for N fertiliser applications (with the EF expressed as a percentage of the total fertiliser N). Country- and practice-specific EFs have been derived for the major emission sources across the different agricultural sectors as described below.

### A1.1 LIVESTOCK HOUSING

#### *Cattle*

Emission factors for two types of cattle housing are currently defined; slurry systems (solid-floor, cubicle housing with scraped passage) and deep litter straw-bedded housing generating farmyard manure (FYM). There is no differentiation between dairy and beef cattle (Table A1.1). The underlying studies from which these EFs are derived are given in Annex 1 (Table AN1).

It is recognised that slatted-floor slurry systems also exist for dairy and beef systems, particularly in Northern Ireland and Scotland, and that the current slurry housing system EF of 27.7% of TAN deposited in the house may not be representative of these systems.

**Table A1.1.** Cattle housing EFs (as % of TAN deposited in the house)

Housing system	EF	SE	n
Slurry, all cattle	27.7	3.85	14
Deep litter (FYM), all cattle	16.8	1.97	10

Seasonal differentiation in the EF is not included in the inventory. The EF for housing might be expected to be greater in summer, because of higher temperatures. However, work by Phillips et al. (1998) showed that summer emissions from dairy cattle housing, where the cattle come in for part of the day for milking, were of a similar magnitude to winter emissions.

#### *Pigs*

As for cattle, housing EFs for pigs have been derived for two management systems, slurry-based and FYM-based, including a larger number of animal categories (Table AN2). A review conducted as part of Defra project AC0123 in 2012 concluded that pig housing has not changed considerably over the inventory reporting period and that the EFs reported here are

relevant for current housing systems. This was largely confirmed (for EF expressed as %TAN) by a more recent housing emissions measurement study funded by AHDB (Dimmock and Stoddart, 2021). However, the latest review into pig housing EFs (commissioned by the EA) led to slight decreases in housing EFs for finishing pigs on slats and weaners on slats (Table A1.2). Most measurements have been made for finishing pigs on either slatted floor or straw-bedded systems, with fewer or no measurements for the other pig categories (Table A1.2).

**Table A1.2.** Pig housing EFs (means and standard errors as % of TAN deposited in the house)

Housing system	EF	SE	n
Dry sows on slats	27.5	9.77*	3
Dry sows on straw	30.8	9.00*	9
Farrowing sows on slats	28.6	2.95*	9
Farrowing sows on straw	33.5		1
Boars on straw	30.8	dry sows value used	
Finishing pigs on slats	28.6	2.11*	19
Finishing pigs on straw	19.6	4.81*	13
Weaners on slats	12.2	4.14*	4
Weaners on straw	7.4		1

\*Differs from previous report due to resolution of inconsistencies

### **Poultry**

Measurements have been made from poultry housing for the poultry categories laying hens, broilers and turkeys (Table AN3). For pullets, breeding hens and other classes of poultry not categorised, a weighted average of the broiler and turkey data were used to derive an emission factor of 14.1% (Table A1.3). Laying hen systems are further categorised as cages (old-style, small battery cages, not permitted after 2012) without belt-cleaning, perchery, free-range and cages (old-style) with belt cleaning, and more modern housing systems as free-range single or multi-tier and colony cages with belt-cleaning (based on data from Defra AC0123).

**Table A1.3.** Poultry housing EFs (means and standard errors as % of TAN deposited in the house)

Housing system	EF	SE	n
Layers, deep pit (* 'old' cages, perchery, free-range)	35.6	8.14	7
*Layers, 'old' cages with belt-cleaning	14.5	4.79	5
Layers free-range single tier	20.1	5.85	3
Layers free-range multi-tier	10.7	3.37	3
Layers colony cages belt-cleaned	8.9	3.15	3
Broilers	9.9	0.76	19
Turkeys	36.2	30.53	3
Pullets, breeding hens and all other poultry	14.1	Based on broilers and turkeys	

\*'old' cages after 2012 are no longer in use, presented because of relevance to historical inventory

### **Sheep**

No specific measurements have been conducted for sheep housing, so the same value is used as for straw-bedded cattle housing i.e. 16.8% of the TAN deposited in the house.

### **Minor livestock**

Horses kept on agricultural holdings have an assumed N excretion of 50 kg per animal per year and are assumed to spend 25% of the year housed. Whereas goats and deer have assumed N excretion of 8.4 and 29.3 kg N per animal per year, respectively and are assumed to spend 8 and 25% of the year housed, respectively. Emission factors (expressed as %TAN) are assumed to be the same as for cattle on FYM.

## **A1.2 HARD STANDINGS (UNROOFED OUTDOOR CONCRETE YARDS)**

### **Cattle**

From Misselbrook et al. (2006), an EF of 75% of the TAN left after scraping is assumed, based on mean measured values of 0.47 and 0.98 g NH<sub>3</sub>-N animal<sup>-1</sup> h<sup>-1</sup> for dairy and beef cattle, respectively, with respective standard errors of 0.09 (n = 28) and 0.39 (n = 30) g NH<sub>3</sub>-N animal<sup>-1</sup> h<sup>-1</sup>.

## **A1.3 MANURE STORAGE**

### **Slurry**

Derived EFs for cattle and pig slurry storage are given in Table A1.4. Measurements from slurry lagoons and above-ground tanks are generally reported as emission per unit area, with only few studies containing sufficient information from which to derive an EF expressed as a percentage of the TAN present in the store (Tables A1.4 and A1.5). The EFs for lagoons are high and substantiated by very little underlying evidence (with no differentiation between pig and cattle slurries) so further measurements are warranted for this source. Currently we assume 4, 10, 24 and 25% of cattle slurry is stored in lagoons for Northern Ireland, Scotland, England and Wales respectively, whereas 4, 24, 25 and 25% of pig slurry is stored in lagoons for Scotland, England, Northern Ireland and Wales respectively. Emissions from below-slat slurry storage inside animal housing are assumed to be included in the animal housing EF, so below-slat storage does not appear as a separate storage category. As only few measurement data are available for EF derivation, and some categories of storage 'read across' from others, a default uncertainty estimate of ± 30% for the 95% confidence interval is suggested for all slurry storage categories.

**Table A1.4.** Slurry storage EF (as % of TAN present in the store)

Storage system	EF	Uncertainty (95% CI)
Cattle slurry above-ground store (no crust)	10 <sup>†</sup>	3.0
Cattle slurry weeping wall	5	1.5
Cattle slurry lagoon (no crust)	52	15.6
Cattle slurry below-ground tank	5 <sup>‡</sup>	1.5
Pig slurry above-ground store	13	3.9
Pig slurry lagoon	52	15.6
Pig slurry below-ground tank	7 <sup>*</sup>	2.1

<sup>†</sup>assumed to be double that of crusted slurry (for which measurements were made); <sup>‡</sup>assumed to be the same as for above-ground slurry store with crust; <sup>\*</sup>assumed to be half the value of above-ground slurry store.

### ***Solid manure***

Derived EFs for cattle, pig and sheep FYM and poultry manure storage are given in Table A1.5. There is large variability in the EF for cattle and pig FYM, with weather conditions in particular influencing emissions, and a combined EF of 28.2% (SE 6.28) is probably justified. Details of the underlying data are given in Tables AN4, AN5 and AN6. The EF for horse FYM is assumed to be the same as that for cattle FYM.

**Table A1.5.** FYM and poultry manure storage EF (as % of TAN present in the store)

Storage system	EF	SE	n
Cattle FYM	26.3	8.28	10
Pig FYM	31.5	10.33	6
Sheep FYM	26.3	Cattle FYM EF used	
Layer manure	14.2	2.99	8
Broiler litter	9.6	2.69	11
Other poultry litter (excluding ducks)	9.6	Broiler litter EF used	
Duck manure	26.3	Cattle FYM EF used	


### **A1.4 MANURE APPLICATION**

Emission factors following manure applications to land are derived using the MANNER\_NPK model (Nicholson et al., 2013), which established standard emission functions using a Michaelis-Menten curve fitting approach for different manure types and applied modifiers according to soil moisture, land use and slurry dry matter content (Table A1.6). Other modifiers included in the model according to wind speed and rainfall within 6 hours of application were not included in the national scale derivation of EF. Modifiers according to application method (splashplate assumed as baseline) and timing of soil incorporation are included as mitigation methods associated with an emission reduction efficiency and are detailed in Appendix 2. Table A1.7 shows the resulting EFs as used in the national inventory. Uncertainties for the weighted average EFs in Table A1.7 were derived from the error terms in the modelled vs. observed plots using the MANNER\_NPK model against UK-specific available data for cattle slurry, pig slurry, FYM (cattle and pig) and poultry manure (Figure A1).

**Table A1.6.** Ammonia EFs and modifiers according to the MANNER\_NPK model

Manure type	Standard EF (as % of TAN applied)	Soil moisture modifier	Land use modifier	Slurry DM modifier	
				Slope	Intercept
Cattle slurry	32.4	x1.3 for dry soil (summer, May-July); x0.7 for moist soil	x0.85 for arable; x1.15 for grassland	8.3	50.2
Pig slurry	25.5	-	-	12.3	50.8
FYM (incl. duck)	68.3	-	-	-	-
Poultry manure	52.3	-	-	-	-

**Figure A1.** MANNER\_NPK model performance against UK data sets for ammonia emissions following land spreading (Nicholson et al., 2013). Cattle slurry (i), pig slurry (ii), FYM (iii) and poultry manure (iv).



Standard errors for the derived slope values were 0.073, 0.148, 0.061 and 0.063 for i, ii, iii and iv, respectively.

**Table A1.7.** Manure application EFs (as % of TAN applied to land)

Manure type	Land use	Season	Slurry DM	EF, %TAN	95% CI, %TAN
Cattle slurry	Grassland	Summer	<4%	32.4	8.4
			4-8%	48.4	
			>8%	64.5	
			Weighted average	52.5	
Cattle slurry	Grassland	Rest of year	<4%	17.4	4.5
			4-8%	26.1	
			>8%	34.7	
			Weighted average	28.2	
Cattle slurry	Arable	Summer	<4%	23.9	6.2
			4-8%	35.8	
			>8%	47.7	
			Weighted average	38.8	
Cattle slurry	Arable	Rest of year	<4%	12.9	3.4
			4-8%	19.3	
			>8%	25.7	
			Weighted average	20.9	
Pig slurry	-	-	<4%	19.2	6.4
			4-8%	31.8	
			>8%	44.3	
			Weighted average	24.2	
FYM (all)	-	-	-	68.3	8.7
Poultry manure (all)	-	-	-	52.3	7.1

## A1.5 GRAZING AND OUTDOOR LIVESTOCK

### ***Cattle and sheep***

The average EF for cattle and sheep (there was no evidence to warrant differentiation) was derived from a number of grazing studies (see Table AN7) with a range of fertiliser N inputs to the grazed pasture. Emissions due to the fertiliser applied to the grazed pasture were discounted using a mean EF for ammonium nitrate applications to grassland (1.4% of N applied). The remaining emission was expressed as a percentage of the estimated urine N (equated here with the TAN in excreta) returned to the pasture by the grazing cattle or sheep. A mean EF of 6% of excreted TAN, with a standard error of 0.7 (n=20) was derived. This value is also assumed for grazing deer and goats.

### ***Outdoor pigs***

Only two studies have made measurements of NH<sub>3</sub> emissions from outdoor pigs (Table AN8), and sufficient data were provided from only one of these to derive a rounded EF of 25% of TAN excreted, with an assumed 95% confidence interval of  $\pm 7.5\%$  of TAN excreted.

### ***Outdoor poultry***

No studies of emissions from outdoor poultry have been reported. An EF of 35% of excreted TAN has been assumed, as it is likely that emissions from freshly dropped excreta will be



substantially lower than from applications of stored manure in which hydrolysis of the uric acid will have occurred to a greater extent. The 95% confidence interval for this EF is assumed to be  $\pm 15\%$  of TAN excreted. A revised 10% of poultry droppings are estimated to be voided outside the house (based on an Environment Agency commissioned review); this is a decrease on the previous estimate.

## A1.6 NITROGEN FERTILISER APPLICATIONS

A model based on Misselbrook et al. (2004) and modified according to data from the Defra-funded NT26 project is used to estimate EFs for different fertiliser types. Each fertiliser type is associated with an  $EF_{max}$  value, which is then modified according to soil, weather and management factors (Table A1.8). EFs are calculated and applied at a 10 km grid resolution, so averaged implied EFs at DA or UK level may vary from year to year. The use of urease inhibitors with urea-based fertilisers and soil placement of N fertiliser are considered as abatement measures and are detailed in Appendix 2.

**Table A1.8.** Nitrogen fertiliser application EF

Fertiliser type	$EF_{max}$ (as % of N applied)	Modifiers <sup>†</sup>
Ammonium nitrate	1.8	None
Ammonium sulphate and diammonium phosphate	45	Soil pH
Urea	45	Application rate, rainfall, temperature
Urea ammonium nitrate	23	Application rate, rainfall, temperature
Other N compounds	1.8	None

### <sup>†</sup>Modifiers:

Soil pH: if calcareous soil, assume EF as for urea; if non-calcareous, assume EF as for ammonium nitrate

Application rate:

- if  $\leq 30$  kg N ha<sup>-1</sup>, apply a modifier of 0.62 to  $EF_{max}$
- if  $\geq 150$  kg N ha<sup>-1</sup>, apply a modifier of 1 to  $EF_{max}$
- if between 30 and 150 kg N ha<sup>-1</sup>, apply a modifier of  $((0.0032 \times \text{rate}) + 0.5238)$

Rainfall: a modifier is applied based on the probability of significant rainfall (>5mm within a 24h period) within 1, 2, 3, 4 or 5 days following application, with respective modifiers of 0.3, 0.5, 0.7, 0.8 and 0.9 applied to  $EF_{max}$ .

Temperature: apply a modifier, with the maximum value constrained to 1, of

$$RF_{temp} = e^{(0.1386 \times (T_{month} - T_{UKannual}))} / 2$$

where  $T_{UKannual}$  is the mean annual air temperature for the UK

An uncertainty bound to the  $EF_{max}$  values of  $\pm 0.3 \times EF_{max}$  is suggested based on the measurements reported under the NT26 project.

## A1.7 DIGESTATE APPLICATIONS TO LAND

### ***Food and crop-based digestates***

Tomlinson et al. (2019) derived an NH<sub>3</sub> EF for surface broadcast digestate (across all types) of 34.7% of the applied N (range 15.4 – 54). Assuming 80% of total N to be in the TAN form, an EF of 43% of TAN applied (range 19 – 68) is derived for use in the agricultural inventory model.

***Livestock manure based digestate***

Literature evidence on the effect of anaerobic digestion on NH<sub>3</sub> emissions at land spreading is mixed, with differing effects of a lower dry matter content (potentially reducing emissions) but higher pH and TAN content (potentially increasing emissions). The assumption applied in the UK inventory is that, expressed as a percentage of the TAN applied, the NH<sub>3</sub> EFs for slurry-digestates are the same as for the corresponding slurry; for cattle and pig FYM-digestates, cattle and pig slurry EFs are applied, respectively, and for poultry manure digestates the value for pig slurry is applied (based on their having similar characteristics).

***Activity data***

Material inputs to anaerobic digestion facilities are derived from the National Non-Food Crops Centre (most recently NNFCC, 2022), with estimated capacity and type of feedstock. Total N content of digestates is based on literature review (Tomlinson et al., 2019) giving mean values of 5.00, 3.97 and 3.35 kg t<sup>-1</sup> for food-waste, energy crop and other organic residue based digestates, respectively, and it is assumed there is no trend across the time series. The TAN content of all digestate types is assumed as 80% of the total N content (RB209).

## ANNEX 1: SOURCES OF UNDERLYING DATA FOR THE UK AMMONIA EMISSION FACTORS

**Table AN1.** Studies delivering cattle housing EFs

Study	Emission g NH <sub>3</sub> -N lu <sup>-1</sup> d <sup>-1</sup>	No. studies	Emission Factor % TAN	Notes on derivation of EF as %TAN
<b>Slurry-based systems</b>				
Demmers et al., 1997	38.6	1	31.1	Dairy cows 1995, assume N excretion of 100 kg N per year
WA0653	21.2	6	19.2	Dairy cows 1998/99, assume N excretion of 105 kg N per year
Dore et al., 2004	72.5	1	53.1	Dairy cows 1998/99, assume N excretion of 105 kg N per year
WAO632/AM110	50.8	3	39.4	Using actual N balance data
Hill, 2000	29.4	1	22.8	Dairy cows 1997, assume N excretion of 104 kg N per year
AM0102	30.5	2	23.7	Dairy cows 2003, assume N excretion of 113 kg N per year
Mean	40.5		31.6	
<b>Weighted mean</b>	<b>34.3</b>		<b>27.7</b>	
<b>Straw-bedded systems</b>				
WA0618 (PT)	20.6	1	18.3	Growing beef, assume N excretion of 56 kg N per year
WAO632/AM110 (PT)	35.0	3	21.6	Using actual N balance data
WA0722	33.2	1	22.9	Dairy cows, 6,500 kg milk per year, therefore assume N excretion of 112 kg N per year
AM0103 (PT)	13.9	1	11.7	Growing beef, values directly from report
AM0103 (Comm farm)	16.7	1	13.4	Dairy cows, assuming 125 g TAN excretion per day (AM0103 report)
AC0102	14.0	3	12.5	Growing beef, assume N excretion of 56 kg N per year
Mean	22.2		16.7	
<b>Weighted mean</b>	<b>23.1</b>		<b>16.8</b>	

**Table AN2. Studies delivering pig housing EFs**

Study	n	Emission factor expressed as:			Av. Live weight	N excretion, kg/place/y
		kg NH <sub>3</sub> /place/y	% TAN	% N		
<b>Dry sows on slats</b>						
Peirson, 1995	2	3.01	22.9	16.0	200	15.5
AHDB, 2021	1	3.65	36.7	25.7	200	11.7
<b>Weighted mean</b>			<b>27.5</b>	<b>19.2</b>		
<b>Dry sows on straw</b>						
Peirson, 1995	2	1.67	12.6	8.9	200	15.5
Koerkamp et al., 1998	1	2.61	19.8	13.9	200	15.5
OC9523	4	4.64	35.3	24.7	200	15.5
AM0102	1 <sup>†</sup>	8.97	68.1	47.7	200	15.5
AHDB, 2021	1	2.29	23.0	16.1	200	11.7
<b>Weighted mean</b>			<b>30.8</b>	<b>21.6</b>		
<b>Farrowing sows on slats</b>						
Peirson, 1995	3	6.46	33.8	23.7	225	22.5
Koerkamp et al., 1998	1	4.41	23.1	16.1	240	22.5
AM0102	3	5.38	30.4	21.3	225	20.8
AHDB, 2021	2	3.76	21.0	14.7	225	21.1
<b>Weighted mean</b>			<b>28.6</b>	<b>20.0</b>		
<b>Farrowing sows on straw</b>						
AHDB, 2021	1	6.01	33.5	23.5	225	21.1
<b>Weaners on slats</b>						
Peirson, 1995	1	0.84	22.5	15.7	12	4.4
Koerkamp et al., 1988	1	0.22	5.9	4.1	12	4.4
AHDB, 2021	2	0.35	10.3	7.2	18	4.0
<b>Weighted mean</b>			<b>12.9</b>	<b>8.6</b>		
<b>Weaners on straw</b>						

AHDB, 2021	1	0.25	7.4	5.1	18	4.0
<b>Finishers on slats</b>						
Peirson, 1995	3	3.18	26.9	18.8	50	13.9
Demmers, 1999	1	2.41	25.3	17.7	25.7	11.2
Koerkamp et al., 1998	1	1.59	16.7	11.7	35	11.2
WA0632	4	3.66	40.4	28.3	60	10.7
WA0720 (fan vent, comm farm)	1	4.59	41.5	29.1	50	13.0
WA0720 (acnv, comm farm)	3	3.42	31.0	21.7	50	13.0
WA0720 (part slat, comm farm)	2	2.28	20.7	14.5	50	13.0
WA0720 (fan vent, Terrington)	1	2.85	21.6	15.2	67.5	15.5
WA0720 (part slat, Terrington)	1	2.31	17.6	12.3	67.5	15.5
AHDB 2016	1	1.87	18.2	12.8	80	12.1
Dimmock, J., Stoddart, H. (2021), AHDB report	1	2.60	26.8	18.8	70	11.4
<b>Weighted mean</b>			<b>28.6</b>	<b>19.4</b>		
<b>Finishers on straw</b>						
Peirson, 1995	2	2.40	20.3	14.2	50	13.9
Koerkamp et al., 1998	1	0.88	9.2	6.4	35	11.2
WA0632	1 <sup>†</sup>	5.65	53.7	37.6	60	12.4
AM0102	1	1.06	9.6	6.7	50	13.0
AM0103 Terrington	2	2.72	23.6	16.7	75	13.4
AM0103 Commercial	1	1.21	10.9	7.7	40	13.0
AC0102	4	1.68	16.6	11.6	45	11.9
AHDB, 2021	1	1.66	17.1	12.0	70	11.4
<b>Weighted mean</b>			<b>19.6</b>	<b>13.7</b>		

<sup>†</sup>Weighting value reduced to 1 from 4 or 5 as values seem to be high outliers

**Table AN3. Studies delivering poultry housing EFs**

Study	Emission g N lu <sup>-1</sup> d <sup>-1</sup>	No. studies	Emission Factor % TAN	Notes
<b>Layers – deep-pit (cages, perchery, free-range)</b>				
Peirson, 1995	79.0	3	22.1	Assume N excretion 0.82 kg (1995)
G Koerkamp, 1998	184.1	1	49.2	Assume N excretion 0.82 kg (1995)
G Koerkamp, 1998	146.1	1	39.0	Assume N excretion 0.82 kg (1995)
WA0368	139.2	1	36.8	Assume N excretion 0.79 kg (1998)
WA0651	196.8	1	57.9	Assume N excretion 0.78 kg (2000)
<b>Weighted mean</b>	<b>129.0*</b>		<b>35.6</b>	
<b>Layers – deep litter: assume same EF as for perchery</b>				
<b>Layers – belt-cleaned (cages)</b>				
Peirson, 1995	36.0	3	10.1	Assume N excretion 0.82 kg (1995)
WA0651 Gleadthorpe	79.2	1	23.3	Assume N excretion 0.78 kg (2000)
WA0651 comm. farm	64.8	1	19.1	Assume N excretion 0.78 kg (2000)
<b>Weighted mean</b>	<b>50.4</b>		<b>14.5</b>	
<b>Layers – Free-range single tier</b>				
AC0123		3	20.1	Refer to AC0123 for details
<b>Layers – Free-range multi-tier</b>				
AC0123		3	10.7	Refer to AC0123 for details
<b>Layers – colony cages with belt cleaning</b>				
AC0123		3	8.9	Refer to AC0123 for details
<b>Broilers</b>				
Demmers et al. 1999	42.0	1	6.7	Assume N excretion 0.56 kg (1995)
Robertson et al 2002	44.0	4	8.3	Assume N excretion 0.55 kg (2000)
Frost et al 2002	54.0	4	9.2	Assume N excretion 0.55 kg (2000)
WA0651 winter	36.0	2	9.5	Derived N excretion from N balance
Mulvenna & Ball AFBI report, 2023	35.1	4	10.1	N excretion derived from Report
WA0651 summer	67.2	2	15.6	Derived N excretion from N balance
WA0651 drinkers	52.8	2	10.9	Derived N excretion from N balance
<b>Weighted mean</b>	<b>46.6</b>		<b>9.9</b>	
<b>Turkeys</b>				
Peirson et al, 1995	92.1*	3	<b>36.2*</b>	Assume N excr 1.59 kg (1995)

lu= livestock unit

\*Value corrected from previous report

**Table AN4. Studies delivering cattle manure storage EFs**

Mean EF g N m <sup>-2</sup> d <sup>-1</sup>	Values g N m <sup>-2</sup> d <sup>-1</sup>	n	Emission as % TAN	Source
<b>Slurry stores and lagoons without crusts</b>				
<b>3.42</b>				Assumed to be double that for crusted stores (WA0641, WA0714)
<b>Slurry stores and lagoons with crusts, weeping wall stores</b>				
<b>1.71</b>	0.6		**2.3	(Phillips et al., 1997)
	1.27, 3.65, 5.7		NA	WA0625
	0.44	2	*6.0	WA0632*
	1.8		NA	WA0641
	1.7		NA	Hill (2000)
	0.48	2	NA	WA0714
	0.5,0.72,0.42,0.73		51.5 (lagoons)	WA0717
	4.2		5.3 (w.wall)	AM0102
			NA	
<b>Below ground slurry tanks</b>				
				Assume same as for crusted above-ground tank
<b>FYM heaps</b>	<b>g N t<sup>-1</sup> initial heap mass</b>			
<b>265</b>	421, 101, 106		NA	WA0618
		2	49	WA0519
		2	29	WA0632
		3	11	Chadwick, 2005
		2	31	WA0716
		1	11	Moral et al., 2012

\*\* Emissions expressed per day. This value assumes 90 d storage.

Slurry stores are assumed to develop a crust unless they are stirred frequently.

Values derived from measurements made using Ferm tubes have been corrected to account for incomplete recovery of ammonia by Ferm tubes (Phillips et al., 1998). (\*IGER values have been corrected using a factor of 0.7).

**Table AN5. Studies delivering pig manure storage EFs**

Mean EF g N m <sup>-2</sup> d <sup>-1</sup>	Values g N m <sup>-2</sup> d <sup>-1</sup>	n	Emission as %TAN	Source
<b>Slurry stores and lagoons</b>				
<b>3.16</b>	1.34	4	13.0	WA0632
	2.47, 6.2		NA	WA0625
	2.4		NA	Phillips et al. (1997)
	1.56		NA	WA0708
	5.0		NA	Phillips et al. (1997)
<b>Below ground slurry tanks</b>				
				Assume 50% of EF for above-ground tank
<b>FYM heaps</b>	<b>g N t<sup>-1</sup> initial heap mass</b>			
<b>1224</b>	539	4	20	WA0632
	1015	2	54	WA0716

Values derived from measurements made using Ferm tubes have been corrected to account for incomplete recovery of ammonia by Ferm tubes (Phillips et al., 1998).

**Table AN6.** Studies delivering poultry manure storage EFs

Mean EF	Values	n	Emission as %TAN	Source
<b>g N t<sup>-1</sup> initial heap mass</b>				
<b>Layer manure</b>				
1956	318	2	3.5	WA0712
	3172	4	14.3	WA0651 (belt scraped)
	3141	1	29.5	WA0651 (deep pit)
	1193	1	20.0	WA0651 (belt scraped)
<b>Litter</b>				
1435	478	1	2.2	WA0712
	1949	4	19.9	WA0651 (winter)
	158	4	1.8	WA0651 (summer)
	639	2	8.4	WA0651 (drinkers)
	3949		NA	WA0716



**Table AN7. Studies delivering cattle and sheep grazing EFs**

	N input	Urine N	NH <sub>3</sub> emission	Due to fertiliser	Due to urine	Emission Factor
	Kg N ha <sup>-1</sup>					%TAN
<b>CATTLE</b>						
<i>Bussink</i>	<i>Fert Res 33 257-265</i>					
1987	550	425	42.2	7.7	34.5	8
1988	550	428	39.2	7.7	31.5	7
1988	250	203	8.1	3.5	4.6	2
<i>Bussink</i>	<i>Fert Res 38 111-121</i>					
1989	250	64.2	3.8	3.5	0.3	0
1989	400	76.2	12.0	5.6	6.4	8
1989	550	94.3	14.7	7.7	7	7
1990	250	217.4	9.1	3.5	5.6	3
1990	400	339	27.0	5.6	21.4	6
1990	550	407.1	32.8	7.7	25.1	6
<i>Lockyer</i>	<i>J Sci Food Agric 35, 837-848</i>					
1	26	0.6455				2
2	26	0.7025				3
<i>Jarvis et al</i>	<i>J Ag Sci 112, 205-216</i>					
1986/87	0	69	6.7	0	6.7	10
1986/87	210	81	9.6	2.94	6.66	8
1986/87	420	207	25.1	5.88	19.22	9
<i>AC0102</i>						
Beef, North Wyke	0			0		10
Beef, Cambridge	0			0		7
<b>SHEEP</b>						
<i>Jarvis et al</i>	<i>J Ag Sci 117, 101-109</i>					
GC	0	169	1.1	0	1.1	1
HN	420	321	8.0	5.88	2.08	1
<i>AC0102</i>						
Boxworth	0					4
North Wyke	0					10

**Table AN8.** Studies delivering EF for outdoor pigs

	Emission g N lu <sup>-1</sup> d <sup>-1</sup>	EF %TAN	Source
Outdoor sows/piglets	25	26.1	Williams et al. (2000)
	66*	NA	Welch (2003)

\*This value is probably an overestimate as emission rates were below the detection limit on a number of occasions (and those data were not included).

The EF was derived from the Williams et al (2000) study, assuming the standard N excretion value for sows and a body weight of 200 kg, giving a mean EF of 25% TAN (assumed to be the same across all animal sub-categories).

## References

- Baines, S., Svoboda, I. F. and Sym, G. (1997) *Estimates of slurry, manure storage and housings in use in Scotland and Northern Ireland*. Report to MAFF (WA0620), SAC Ayr.
- Bussink, D.W. (1994). Relationship between ammonia volatilization and nitrogen fertilizer application rate, intake and excretion of herbage nitrogen by cattle on grazed swards. *Fertilizer Research* **38**, 111-121
- CAMAR: Groot Koerkamp, P. W. G., Metz, J. H. M., Uenk, G. H., Phillips, V. R., Holden, M. R., Sneath, R. W., Short, J. L., White, R. P., Hartung, J., Seedorf, J., Schröder, M., Linkert, K. H., Pedersen, S., Takai, H., Johnsen, J. O. and Wathes, C. M., 1998. Concentrations and emissions of ammonia in livestock buildings in Northern Europe. *Journal of Agricultural Engineering Research* **70**, 79-95.
- Chalmers, A. G., et al. (2001). *Fertiliser use on farm crops for crop year 2000*. British Survey of Fertiliser Practice, Edinburgh: The Stationery Office.
- Chadwick, D.R. (2005). Emissions of ammonia, nitrous oxide and methane from cattle manure heaps: effect of compaction and covering. *Atmospheric Environment* **39**, 787-799.
- Chambers, B.J., Smith, K.A. and van der Weerden, T.J. (1997). Ammonia emissions following the land spreading of solid manures. In *Gaseous Nitrogen Emissions from Grasslands*. Eds S.C. Jarvis and B.F. Pain, CAB International, Oxford, pp. 275-280.
- Chambers, B. J., Lord, E. I., Nicholson, F. A. and Smith, K. A. (1999). Predicting nitrogen availability and losses following application of organic manures to arable land: MANNER. *Soil Use and Management* **15**, 137-143.
- Cumby T., Sandars D., Nigro E., Sneath R. and Johnson G. (2005) Physical assessment of the environmental impacts of centralised anaerobic digestion. Report by Silsoe Research Institute. 112pp.
- DANI (1998). Statistical Review of Northern Ireland Agriculture, 1997. Department of Agriculture for Northern Ireland, Economics and Statistics Division, Belfast, Northern Ireland, UK.
- Defra (2001) [http://www.defra.gov.uk/esg/work\\_hm/publications/cs/fps/fpsfinalreport.PDF](http://www.defra.gov.uk/esg/work_hm/publications/cs/fps/fpsfinalreport.PDF)



- Demmers, T.G.M., Phillips, V.R., Short, J.L., Burgess, L.R., Hoxer, R.P. and Wathes, C.M (1997). Validation of ventilation rate measurement methods and the ammonia emission from a naturally-ventilated UK dairy and beef unit. In: *Ammonia and Odour Emissions from Animal Production Facilities*. Eds J.A.M. Voermans and G.J. Montenevy, Proceedings of an international symposium held at Vinkeloord, Netherlands, 6-10 October 1997. Published by NTVL, Rosmalen, NL pp. 219-230.
- Demmers, T.G.M., Burgess, L.R., Short, J.L., Phillips, V.R., Clark, J.A. and Wathes, C.M. (1999). Ammonia emissions from two mechanically ventilated UK livestock buildings. *Atmospheric Environment* **33**, 217-227.
- Dimmock, J., Stoddart, H. (2021). New Ammonia Emission Factors for the English Pork Industry. AHDB Report, March 2021.
- Dore, C. J., Jones, B. M. R., Scholtens, R., Burgess, L. R., Huis in't Veld, J. W. H., Phillips, V. R. (2004). Robust methods for measuring ammonia emission rates from livestock buildings and manure stores. Part 1 Comparative demonstrations of three methods on the farm. *Atmospheric Environment* **38**, 3017-3024.
- Dragosits, U., Jones S.K., Vogt E. and Sutton M.S. (2006) 2005 Update on Ammonia emissions from non-agricultural sources for the NAEI. CEH Report AS06/20. Centre for Ecology & Hydrology Edinburgh, Bush Estate, Penicuik. 14pp.
- Elliott, M., Ingeldew, D., Richmond, B., Del Vento, S., Gorji, S., Hamilton, N., Karagianni, E., Kelsall, A., Pang, Y., Passant, N., Pearson, B., Richardson, J., Stewart, R., Thistlethwaite, G., Tsagatakis, I., Wakeling, D., Wiltshire, J., Wong, J., Hobson, M., Gibbs, M., Dore, C., Thornton, A., Anthony, S., Carswell, A., Gilhespy, S., Cardenas, L., Dragosits, U., Tomlinson, S. (2024). UK Informative Inventory Report (1990 to 2022). URL: [https://naei.beis.gov.uk/reports/reports?report\\_id=1135](https://naei.beis.gov.uk/reports/reports?report_id=1135)
- Hill, R.A. (2000). Emission, dispersion and local deposition of ammonia volatilised from farm buildings and following the application of cattle slurry to grassland. PhD Thesis, University of Plymouth.
- Hodge, I. and Renwick, A. (2006). Business as usual projections of agricultural activities for the water framework directive: Phase 2. Final Report. Rural Business Unit, Environmental Economy and Policy Research Group, Department of Land Economy, 19 Silver Street, Cambridge CB3 9EP.
- Jarvis, S.C; Hatch, D. J; Orr, R.J. and Reynolds, S.E. (1991). Micrometeorological studies of ammonia emissions from sheep grazed swards. *Journal of Agricultural Science Cambridge*, **117**, 101-109
- Jarvis, S. C. and Bussink, D. W. (1990). Nitrogen losses from grazed swards by ammonia volatilization. Proceedings of the 13th General Meeting of the European Grassland Federation, June 25-29, 1990, Banska Bystrica, Czechoslovakia, p.13-17.
- Kirchmann, H., and Witter, E. (1989). Ammonia volatilization during aerobic and anaerobic manure decomposition. *Plant and Soil* **115**, 35-41.
- Koerkamp, P., Metz, J. H. M., Uenk, G. H., Phillips, V. R., Holden, M. R., Sneath, R. W., Short, J. L., White, R. P., Hartung, J., Seedorf, J., Schroder, M., Linkert, K. H., Pedersen, S., Takai, H.,

- Johnsen, J. O. and Wathes, C. M. (1998). Concentrations and emissions of ammonia in livestock buildings in Northern Europe. *Journal of Agricultural Engineering Research* **70**, 79-95.
- Ledgard, S. F. (1996). Nitrogen inputs and losses from New Zealand dairy farmlets, as affected by nitrogen fertilizer applications: year one. *Plant and Soil* **181**, 65-69.
- MAFF (2000). *Fertilizer Recommendations for Agricultural and Horticultural Crops (RB209)*, 7<sup>th</sup> Edition. Her Majesty's Stationery Office, London, UK.
- Mercer, D. R. (1993) *Estimates of the number and types of poultry housing in use in England and Wales*. Report to MAFF, ADAS Nottingham.
- Misselbrook, T.H., Webb, J. and Gilhespy, S.L. (2006). Ammonia emissions from outdoor concrete yards used by livestock – quantification and mitigation. *Atmospheric Environment* **40**, 6752-6763.
- Misselbrook, T.H., Sutton, M.A. and Scholefield, D. (2004). A simple process-based model for estimating ammonia emissions from agricultural land after fertilizer applications. *Soil Use and Management* **20**, 365-372.
- Misselbrook, T. H., Smith, K. A., Johnson, R. A. and Pain, B. F. (2002). Slurry application techniques to reduce ammonia emissions: Results of some UK field-scale experiments. *Biosystems Engineering* **81**, 313-321.
- Misselbrook, T. H., Webb, J., Chadwick, D. R., Ellis, S. and Pain, B. F. (2001). Gaseous emissions from outdoor concrete yards used by livestock. *Atmospheric Environment* **35**, 5331-5338.
- Misselbrook, T. H., Pain, B. F. and Headon, D. M. (1998). Estimates of ammonia emission from dairy cow collecting yards. *Journal of Agricultural Engineering Research* **71**, 127-135.
- Moral, R., Bustamante, M.A., Chadwick, D.R., Camp, V., Misselbrook, T.H., 2012. N and C transformations in stored cattle farmyard manure, including direct estimates of N-2 emission. *Resources Conservation and Recycling* **63**, 35-42.
- Mulvenna, C.C. and Ball, M.E.E. (2023). The ammonia emissions from mechanically ventilated broiler houses in Northern Ireland – Updated figures 2023. Agri-Food & Biosciences Institute, Department of Agriculture, Environment and Rural Affairs evidence and Innovation Project 17/04/03 (48168) – Evaluation of ammonia and greenhouse gas emissions from the livestock production sector in Northern Ireland
- NNFCC (2020) Anaerobic Digestion Deployment in the United Kingdom (<http://www.nnfcc.co.uk/publications/report-anaerobic-digestion-deployment-in-the-uk>). (Accessed August 2020).
- Nicholson, F.A., Bhogal, A., Chadwick, D., Gill, E., Gooday, R.D., Lord, E., Misselbrook, T., Rollett, A.J., Sagoo, E., Smith, K.A., Thorman, R.E., Williams, J.R., Chambers, B.J. (2013). An enhanced software tool to support better use of manure nutrients: MANNER-NPK. *Soil Use and Management* **29**, 473-484.
- Nicholson F., Bhogal A., Cardenas L., Chadwick D., Misselbrook T., Rollett A., Taylor M., Thorman R., and Williams J. (2017) Nitrogen losses to the environment following food-based digestate and compost applications to agricultural land. *Environmental Pollution* **228**, 504-516.



- Nicholson, F. A., Chambers, B. J. and Smith, K. A. (1996) Nutrient composition of poultry manures in England and Wales. *Bioresource Technology* **58**, 279-284.
- Nicholson, R. J. and Brewer, A. J. (1994) *Estimates of the numbers and types of slurry and manure stores in use in England and Wales related to livestock species*. Report to DEFRA (WA0611), ADAS Cambridge.
- Pain, B. F., Rees, Y. J. and Lockyer, D. R. (1988). Odour and ammonia emission following the application of pig or cattle slurry to land. In: *Volatile emissions from livestock farming and sewage operations*, eds V C Neilsen, J H Voorburg and P L'Hermite. Elsevier Applied Science, London, pp. 2 - 11.
- Pain, B. F., Phillips, V. R., Clarkson, C. R. and Klarenbeek, J. V. (1989). Loss of nitrogen through ammonia volatilisation following the application of pig or cattle slurry to grassland. *Journal of the Science of Food and Agriculture* **47**, 1-12.
- Peirson, S. (1995). Measurement of odour and ammonia emissions from livestock buildings, Phase 1 - Final Report to MAFF. Project no. WAO601, ADAS Beverley.
- Phillips, V.R., Sneath, R.W., Williams, A.G., Welch, S.K., Burgess, L.R., Demmers, T.G.M. and Short, J.L. (1997). Measuring emission rates of ammonia, methane and nitrous oxide from full-sized slurry and manure stores. In: *Ammonia and Odour Emissions from Animal Production Facilities*. Eds J.A.M. Voermans and G.J. Montaney, Proceedings of an international symposium held at Vinkeloord, Netherlands, 6-10 October 1997. Published by NTVL, Rosmalen, NL pp. 197-208.
- Phillips, V. R., Bishop, S. J., Price, J. S. and You, S. (1998). Summer emissions of ammonia from a slurry-based, UK, dairy cow house. *Bioresource Technology* **65**, 213-219.
- Robertson, A. P., Hoxey, R. P., Demmers, T. G. M., Welch, AS. K., Sneath, R. W., Stacey, K. F., Fothergill, A., Filmer, D. and Fisher, C. (2002). Commercial-scale studies of the effect of broiler-protein intake on aerial pollutant emissions. *Biosystems Engineering* **82**, 217-225.
- Sheppard, A. (1998) The Structure of Pig Production in England and Wales. Results of the National Survey of Pig Production Systems. *Special Studies in Agricultural Economics Report No. 40*, University of Exeter.
- Smith, K. A. and Chambers, B. J. (1995). Muck from waste to resource - utilization: the impacts and implications. *Agricultural Engineer*, **50**, 33-38.
- Smith, K. A. and Frost, J. P. (2000). Nitrogen excretion by farm livestock with respect to land spreading requirements and controlling nitrogen losses to ground and surface waters. Part 1: cattle and sheep. *Bioresource Technology* **71**, 173-181.
- Smith, K. A., Charles, D. R. and Moorhouse, D. (2000a). Nitrogen excretion by farm livestock with respect to land spreading requirements and controlling nitrogen losses to ground and surface waters. Part 2: pigs and poultry. *Bioresource Technology* **71**, 183-194.
- Smith, K. A., Jackson, D. R., Misselbrook, T. H., Pain, B. F. and Johnson, R. A. (2000b). Reduction of ammonia emission by slurry application techniques. *Journal of Agricultural Engineering Research* **77**, 277-287.

- Smith, K. A., Brewer, A. J., Dauven, A. and Wilson, D. W. (2000c). A survey of the production and use of animal manures in England and Wales. I. Pig manure. *Soil Use and Management* **16**, 124-132.
- Smith, K. A., Brewer, A. J., Crabb, J. and Dauven, A. (2001a). A survey of the production and use of animal manures in England and Wales. II. Poultry manure. *Soil Use and Management* **17**, 48-56.
- Smith, K. A., Brewer, A. J., Crabb, J. and Dauven, A. (2001b). A survey of the production and use of animal manures in England and Wales. III. Cattle manures. *Soil Use and Management* **17**, 77-87.
- Smith, K. (2012) Manure management. Report as part of Defra AC0114 (including Appendix).
- Sommer, S.G., Christensen, B.T., Nielsen, N.E., and Schjrrring, J.K. (1993). Ammonia volatilization during storage of cattle and pig slurry: effect of surface cover. *Journal of Agricultural Science, Cambridge*, **121**, 63-71.
- Sutton, M. A., Dragosits, U., Tang, Y. S. and Fowler, D. (2000). Ammonia emissions from non-agricultural sources in the UK. *Atmospheric Environment* **34**, 855-869.
- Thompson, R. B., Pain, B. F. and Lockyer, D. R. (1990a). Ammonia volatilization from cattle slurry following surface application to grassland. I. Influence of mechanical separation, changes in chemical composition during volatilization and the presence of the grass sward, *Plant and Soil* **125**, 109-117.
- Thompson, R. B., Pain, B. F. and Rees, Y. J. (1990b). Ammonia volatilization from cattle slurry following surface application to grassland. II. Influence of application rate, windspeed and applying slurry in narrow bands. *Plant and Soil* **125**, 119-128.
- Tomlinson S.J., Thomas I.N., Carnell E.J., and Dragosits U. (2019). Reviewing estimates of UK ammonia emissions from landfill, composting & anaerobic digestion: Improvement Plan 2018. Report for Defra (AQ\_IP\_2018\_20). April 2019. 63pp.
- van der Weerden, T. J. and Jarvis, S. C. (1997). Ammonia emission factors for N fertilisers applied to two contrasting grassland soils. *Environmental Pollution* **95**, 205-211.
- Wathes, C. M., Holden, M. R., Sneath, R. W., White, R. P. and Phillips, V. R. (1997). Concentrations and emission rates of ammonia, nitrous oxide, methane, carbon dioxide, dust and endotoxin in UK broiler and layer houses. *British Poultry Science* **38**, 14-28.
- Webb, J., (2001). Estimating the potential for ammonia emissions from livestock excreta and manures. *Environmental Pollution* **111**, 395-406.
- Webb, J., and Misselbrook, T. H. (2004). A mass-flow model of ammonia emissions from UK livestock production. *Atmospheric Environment* **38**, 2163-2176.
- Webb, J., Misselbrook, T., Pain, B. F., Crabb, J. and Ellis, S. (2001). An estimate of the contribution of outdoor concrete yards used by livestock to the UK inventories of ammonia, nitrous oxide and methane. *Atmospheric Environment* **35**, 6447-6451.
- Welch, D.C. (2003) A methodology for the measurement of distributed agricultural sources of ammonia outdoors. PhD thesis, University of Nottingham.

Williams, J. R., Chambers, B. J., Hartley, A. R., Ellis, S. and Guise, H. J. (2000). Nitrogen losses from outdoor pig farming systems. *Soil Use and Management* **16**, 237-243.

WRAP 2014. A survey of the UK Anaerobic Digestion industry in 2013. Waste and Resources Action Programme. <http://www.wrap.org.uk/content/survey-uk-anaerobic-digestion-industry-2013>. (Accessed September 2016).

WRAP 2016a. Field Experiments for Quality Digestate and Compost in Agriculture. Waste and Resources Action Programme. <http://www.wrap.org.uk/content/digestate-and-compost-agriculture-dc-agri-reports>. (Accessed September 2016).

### DEFRA Projects

Final reports from the following projects are available from Defra:

AC0114	GHG Platform – data management
AC0123	Developing new ammonia emission factors for modern livestock housing (Phase 2)
AM0101	National ammonia reduction strategy evaluation system (NARSES)
AM0102	Modelling and measurement of ammonia emissions from ammonia mitigation pilot farms
AM0103	Evaluation of targeted or additional straw use as a means of reducing ammonia emissions from buildings for housing pigs and cattle
AM0110	Additional housing measurements for solid vs. liquid manure management systems
AM0111	Measurement and abatement of ammonia emissions from hard standings used by livestock
AM0115	Investigation of how ammonia emissions from buildings housing cattle vary with the time cattle spend inside them
DO108	Food and Agriculture Policy Research Institute – UK Project
ES0116	Field work to validate the manure incorporation volatilization system (MAVIS)
KT0105	Manure Nutrient Evaluation Routine (MANNER-NPK)
LK0643	UK Poultry Industry IPPC Compliance (UPIC)
NT2001	Integration of animal manures in crop and livestock farming systems: nutrient demonstration farms
NT2402	Impact of nutrition and management on N and P excretions by dairy cows
NT2605	The behaviour of some different fertiliser-N materials - Main experiments
OC9117	Ammonia emission and deposition from livestock production systems
WA0519	Enhancing the effective utilisation of animal manures on-farm through effective compost technology
WA0618	Emissions from farm yard manure based systems for cattle
WA0625	The effects of covering slurry stores on emissions of ammonia, methane and nitrous oxide
WA0632	Ammonia fluxes within solid and liquid manure management systems
WA0633	Predicting ammonia loss following the application of organic manures to land
WA0638	Low cost, aerobic stabilisation of poultry layer manure
WA0641	Low-cost covers to abate gaseous emissions from slurry stores

WA0651	Ammonia fluxes within broiler litter and layer manure management systems
WA0652	Field ammonia losses in sustainable livestock LINK Project LK0613
WA0653	Quantifying the contribution of ammonia loss from housed dairy cows to total N losses from dairy systems (MIDaS2)
WA0707	Effect of storage conditions on FYM composition, gaseous emissions and nutrient leaching during storage
WA0708	Covering a farm scale lagoon of pig slurry
WA0712	Management techniques to minimise ammonia emissions during storage and land spreading of poultry manures
WA0714	Natural crusting of slurry storage as an abatement measure for ammonia emission on dairy farms
WA0716	Management techniques to reduce ammonia emissions from solid manures
WA0717	Ammonia emissions and nutrient balance in weeping-wall stores and earth banked lagoons for cattle slurry storage
WA0720	Demonstrating opportunities of reducing ammonia emissions from pig housing
WA0722	Ammonia emission from housed dairy cows in relation to housing system and level of production
WT0715NVZ	Nitrogen and phosphorus output standards for farm livestock



## APPENDIX 2: REDUCTION EFFICIENCIES FOR AMMONIA MITIGATION METHODS APPLICABLE TO THE UK AMMONIA EMISSION INVENTORY

### INTRODUCTION

Agriculture is the major source of ammonia (NH<sub>3</sub>) emissions to the atmosphere in the UK, accounting for >80% of anthropogenic emissions. Most of these emissions derive from urea excreted by farmed livestock (or uric acid in the case of poultry) and emissions will therefore arise wherever livestock excreta are deposited or managed i.e. at grazing, in livestock housing and during manure storage and application to land. Emissions also arise from inorganic nitrogen (N) fertilisers applied to land. The emission factors used to quantify these emissions in the national inventory are reported separately. A growing number of potential mitigation methods applicable to one or more of the emission sources have been described in the literature. This report lists those that are currently included in the inventory of NH<sub>3</sub> emissions from UK agriculture together with the mean NH<sub>3</sub> emission reduction efficiency associated with each method. In addition, the current state of knowledge regarding the impact of the implementation of each method on emissions of nitrous oxide and methane is given so that these mitigation methods can be fully included in the revised combined agricultural greenhouse gas (GHG) and NH<sub>3</sub> emission inventory.

### Emission reduction methods

Only explicit mitigation methods are included here – i.e. those that are associated with a reduction in the emission factor for a particular source. Implicit mitigation methods, generally associated with efficiency improvements (e.g. a reduction in N fertiliser use through better accounting for manure N use; a reduction in livestock numbers associated with productivity improvements), will be reflected in the inventory through changes in the activity data and are not described here. One exception in the current NH<sub>3</sub> emission inventory is the inclusion of a dietary measure, namely low crude protein diets for dairy cows, which is associated with a 20% reduction in the ammoniacal N content of dairy cow excreta over the housed winter period. In the revised emission inventories, N excretion will be derived using a balance approach according to diet and production characteristics and will therefore reflect any changes in the crude protein content of the diet.

Mitigation methods are categorised according to the emission source i.e. livestock housing, hard standings, manure storage, manure spreading and fertiliser application. Data sources are given, but the reported emission reduction efficiencies are not necessarily the arithmetic mean of reported studies but are more aligned with the expert judgement approaches used in the Defra 'Mitigation Methods - User Guide' (Newell Price et al., 2011) and the UNECE Task Force for Reactive Nitrogen 'Options for Ammonia Mitigation Guidance Document' (Bittman

et al., 2014). These documents and other cited literature should be consulted for more detailed information on the mitigation methods included in Table A2.1.

Uncertainties are not well defined for these emission reduction estimates, so following 2006 IPCC Guidelines for Tier 2 approach to estimating emissions from manure management, uncertainty bound of  $\pm 20\%$  of the reported value are applied with constraining limits of 0 and 100% also implemented.

**Table A2.1.** Reduction efficiencies for ammonia emission mitigation methods and an indication of their impact on nitrous oxide and methane emissions

Emission source	Mitigation method	Ammonia emission reduction efficiency (%)	Nitrous oxide <sup>†</sup>	Methane <sup>†</sup>	Data source
Cattle housing	Increased scraping frequency in cubicle house (from 2 to 4x per day)	15	-	-	Webb et al. (2006); Braam et al. (1997)
	Grooved flooring system for rapid urine draining	35	-	-	Swiestra et al. (2001); Bittman et al. (2014)
Pig housing	Partly slatted floor with reduced pit area	30	-	-	Bittman et al. (2014)
	Acid air scrubbing techniques	80	-	-	Bittman et al. (2014)
	Frequent slurry removal with vacuum system	25	-	-	Bittman et al. (2014)
	Floating balls on below-slat slurry surface	25	-	-	Bittman et al. (2014)
Poultry housing	Air drying of manure on laying hen manure belt systems	30	?	?	Bittman et al. (2014)
	Acid air scrubbing techniques	80	-	-	Bittman et al. (2014)
	Poultry litter drying	60	?	?	Defra WA0638; Defra AC0123
Dairy cow collecting yards	Wash down with water twice per day	70	-	-	Misselbrook et al. (2006)
Slurry storage	Crusting of cattle slurry	50	☒ EF from 0 to 0.005 (IPCC 2006)	↓ Methane Conversion Factor from 17 to 10% (IPCC 2006)	Misselbrook et al. (2005)

	Floating cover (e.g. expanded clay granules)	60	-	-	Bittman et al. (2014); Defra AC0115
	Tight lid, roof or tent structure	80	-	-	Bittman et al. (2014)
FYM/poultry manure storage	Sheeting cover	60	↓ by 30%	-	Chadwick (2005)
Slurry application	Trailing hose	30	-	-	Smith et al. (2000); Misselbrook et al. (2002); Bittman et al. (2014)
	Trailing shoe	60	-	-	Smith et al. (2000); Misselbrook et al. (2002); Bittman et al. (2014)
	Shallow injection	70	-	-	Smith et al. (2000); Misselbrook et al. (2002); Bittman et al. (2014)
Cattle slurry to arable	Incorporation within 4h by plough	59	-	-	Defra ES0116
	Incorporation within 4h by disc	52	-	-	Defra ES0116
	Incorporation within 4h by tine	46	-	-	Defra ES0116
	Incorporation within 24h by plough	21	-	-	Defra ES0116
	Incorporation within 24h by disc	19	-	-	Defra ES0116
	Incorporation within 24h by tine	17	-	-	Defra ES0116
Pig slurry to arable	Incorporation within 4h by plough	67	-	-	Defra ES0116
	Incorporation within 4h by disc	59	-	-	Defra ES0116
	Incorporation within 4h by tine	52	-	-	Defra ES0116
	Incorporation within 24h by plough	29	-	-	Defra ES0116
	Incorporation within 24h by disc	26	-	-	Defra ES0116
	Incorporation within 24h by tine	23	-	-	Defra ES0116
Cattle, pig and duck FYM	Incorporation within 4h by plough	71	-	-	Defra ES0116

	Incorporation within 4h by disc	55	-	-	Defra ES0116
	Incorporation within 4h by tine	24	-	-	Defra ES0116
	Incorporation within 24h by plough	34	-	-	Defra ES0116
	Incorporation within 24h by disc	27	-	-	Defra ES0116
	Incorporation within 24h by tine	11	-	-	Defra ES0116
Poultry manure	Incorporation within 4h by plough	86	-	-	Defra ES0116
	Incorporation within 4h by disc	73	-	-	Defra ES0116
	Incorporation within 4h by tine	64	-	-	Defra ES0116
	Incorporation within 24h by plough	60	-	-	Defra ES0116
	Incorporation within 24h by disc	50	-	-	Defra ES0116
	Incorporation within 24h by tine	44	-	-	Defra ES0116
Urea fertiliser	Urease inhibitor	70	↓ (Smith et al. 2012)	-	Defra NT26
UAN fertiliser	Urease inhibitor	40	?	-	Defra NT26

† □ increase in emission; ↓ decrease in emission; - no effect; ? uncertain of effect

## References

- Bittman, S., Dedina, M., Howard, C.M., Oenema, O., Sutton, M.A., (eds), 2014. *Options for Ammonia Mitigation: Guidance from the UNECE Task Force on Reactive Nitrogen*, Centre for Ecology and Hydrology, Edinburgh, UK
- Newell Price et al 2011 User Guide
- Braam, C.R., Ketelaars, J., Smits, M.C.J., 1997. Effects of floor design and floor cleaning on ammonia emission from cubicle houses for dairy cows. *Netherlands Journal of Agricultural Science* 45, 49-64.
- Defra AC0115 Improvements to the national inventory: Methane
- Defra AC0123 Developing new ammonia emission factors for modern livestock housing (Phase 2)
- Defra ES0116 Field work to validate the manure incorporation volatilization system (MAVIS)
- Defra WA0638 Lost cost, aerobic stabilisation of poultry layer manure
- Misselbrook, T.H., Brookman, S.K.E., Smith, K.A., Cumby, T.R., Williams, A.G., McCrory, D.F., 2005. Crusting of stored dairy slurry to abate ammonia emissions: pilot-scale studies. *Journal of Environmental Quality* 34, 411-419.
- Misselbrook, T.H., Smith, K.A., Johnson, R.A., Pain, B.F., 2002. Slurry application techniques to reduce ammonia emissions: Results of some UK field-scale experiments. *Biosystems Engineering* 81, 313-321.
- Misselbrook, T.H., Webb, J., Gilhespy, S.L., 2006. Ammonia emissions from outdoor concrete yards used by livestock - quantification and mitigation. *Atmospheric Environment* 40, 6752-6763.
- Smith, K.A., Dobbie, K.E., Thorman, R., Watson, C.J., Chadwick, D.R., Yamulki, S., Ball, B.C., 2012. The effect of N fertilizer forms on nitrous oxide emissions from UK arable land and grassland. *Nutrient Cycling in Agroecosystems* 93, 127-149.
- Smith, K.A., Jackson, D.R., Misselbrook, T.H., Pain, B.F., Johnson, R.A., 2000. Reduction of ammonia emission by slurry application techniques. *Journal of Agricultural Engineering Research* 77, 277-287.
- Swierstra, D., Braam, C.R., Smits, M.C., 2001. Grooved floor system for cattle housing: Ammonia emission reduction and good slip resistance. *Applied Engineering in Agriculture* 17, 85-90.
- Webb, J., Ryan, M., Anthony, S.G., Brewer, A., Laws, J., Aller, M.F., Misselbrook, T.H., 2006. Cost-effective means of reducing ammonia emissions from UK agriculture using the NARSES model. *Atmospheric Environment* 40, 7222-7233.