



Air Quality Pollutant Inventories for England, Scotland, Wales and Northern Ireland: 1990 – 2010

A report of the National Atmospheric Emissions Inventory

September, 2012

Air Quality Pollutant Inventories for England, Scotland, Wales and Northern Ireland: 1990 – 2010

Main authors

J MacCarthy, G Thistlethwaite, E Salisbury, Y Pang,
T Misselbrook

September, 2012

Title	Air Quality Pollutant Inventories, for England, Scotland, Wales and Northern Ireland: 1990 – 2010	
Customers	Department for Environment, Food and Rural Affairs, The Scottish Government, The Welsh Government, The Northern Ireland Department of Environment.	
Confidentiality, copyright reproduction	and	Crown Copyright
NAEI reference	56186/2010/CD7720/JMC	
ISBN	978-0-9573549-0-6	
Report number	AEA/ENV/R/3323	
Issue number	Final 1.0	

AEA Group
The Gemini Building
Fermi Avenue
Didcot
Oxfordshire
OX11 0QR

Telephone 0870 190 6584

AEA is a business name of AEA Technology plc
AEA is certified to ISO9001 and ISO14001

Main Authors	AEA: J. MacCarthy, G. Thistlethwaite, Y. Pang Aether: E. Salisbury Rothamsted Research: T. Misselbrook
---------------------	--

Approved by	Name	Justin Goodwin
	Signature	
	Date	12 th September, 2012

Executive Summary

This is the Air Quality Pollutant Inventory Report for England, Scotland, Wales and Northern Ireland. The report presents emission inventories for the constituent countries of the UK for the period 1990 to 2010, for the following priority Air Quality (AQ) pollutants:

- Ammonia (NH₃)
- Carbon monoxide (CO)
- Nitrogen oxides (NO_x as NO₂)
- Non-methane volatile organic compounds (NMVOCs)
- Sub-10 micron particulate matter (PM₁₀)
- Sulphur dioxide (SO₂)
- Lead (Pb)

These inventories are compiled on behalf of the UK Department for Environment, Food & Rural Affairs, the Scottish Government, the Welsh Government and the Department of Environment for Northern Ireland, by the UK emission inventory teams at AEA, Aether and Rothamsted Research.

Data Sources and Inventory Methodology

The constituent country inventories are compiled by disaggregating the UK emission totals presented within “UK Informative Inventory Report 1980 to 2010” (Passant *et al.*, 2012), derived from the National Atmospheric Emissions Inventory (NAEI) database. The emission estimates for each pollutant are presented in NFR format, to be consistent with the UK inventory submissions to the United Nations Economic Commission for Europe (UNECE), which follow international inventory reporting guidelines.

The method for disaggregating UK emission totals across the constituent countries draws on a combination of point source data (e.g. Pollution Inventory¹ data for industrial emissions) and sub-national and local datasets such as:

- DECC sub-national statistics on energy use
- Other regional energy use data for specific industries or regional data on raw material consumption or sector-specific production
- Major road traffic count data
- Domestic and international flight data for all major UK airports
- Rail company fuel use estimates
- Regional housing, employment, population and consumption data
- Agricultural surveys (livestock numbers, crop production, fertiliser application)
- Land use survey data

Emissions from the offshore oil & gas exploration and production sector are not attributed to a specific country inventory, but are reported within an “unallocated” category. Note, however, that emissions from onshore oil & gas terminals are assigned to the appropriate country inventories.

For many sources of AQ pollutants, the data available for constituent country emissions are less detailed than for the UK as a whole, and for some sources, country-level data are not available at all. In particular, detailed energy balances to provide annual fuel-specific consumption data by source sector are not available for England, Scotland, Wales and Northern Ireland.

Sub-national energy statistics are published annually by the Department for Energy and Climate Change (DECC) within the quarterly Energy Trends² publication. These statistics are limited in their detail when

¹The term “Pollution Inventory” is used here to represent the industrial emissions databases of the UK environmental regulators (The Environment Agency of England & Wales, the Scottish Environment Protection Agency and the Northern Ireland Department of Environment), which comprise annual emission estimates from all IPC/IPPC-regulated processes under their authority.

compared to UK-level energy statistics, but do provide estimated fuel use data for England, Scotland, Wales and Northern Ireland for the following source sectors:

- Industry (1A2) & Commercial (1A4a)
- Agriculture (combustion sources) (1A4c)
- Residential (1A4b)

The DECC sub-national energy statistics have been developed in recent years to provide estimates of fuel use and CO₂ emissions data at Local Authority (LA) level across the UK. The latest available data include LA solid and liquid fuel use estimates for 2005 to 2009, with gas and electricity data also being available up to 2010.

The DECC data at local and regional level are derived from analysis of gas and electricity meter point data, supplemented by additional research to estimate the distribution of solid fuels and petroleum-based fuels across the UK. Since the initial study and presentation of experimental data for 2003 and 2004, each annual revision to the local and regional data has included data improvements through targeted sector research. These DECC sub-national energy statistics continue to evolve and improve, reducing data inaccuracies and are the best data available to inform the patterns of fuel use across the Devolved Administrations. They are therefore used to underpin the pollutant emission estimates from fuel combustion sources within the inventories presented here, in conjunction with other data sources such as EU Emissions Trading System (EU ETS) fuel use data for large industrial sites and other DA-specific energy data.

For other significant emission sources there are complete country-level datasets available, although some of these are less detailed than data used for the UK Inventory:

- Industrial process emissions are based on plant operator estimates reported to environmental agencies under regulatory systems such as Integrated Pollution Prevention and Control (IPPC). Major sources include power stations, cement and lime kilns, iron & steelworks, aluminium and other non-ferrous metal plant, chemical industries. These data are not available across the full time-series from 1990, as the regulatory reporting regimes developed in the late 1990s (in England, Wales and Northern Ireland) and early 2000s (in Scotland);
- Emissions from oil and gas terminals and offshore platforms and rigs, are based on operator estimates reported to the DECC Oil and Gas team in Aberdeen through the Environmental Emissions Monitoring System, EEMS;
- Agricultural emissions are based on UK emission factors and annual survey data across each of the Devolved Administrations, including estimates of arable production and livestock numbers;
- Emissions from waste disposal activities are estimated based on modelled emissions from the UK air quality inventory (Defra, 2012) split out across the DAs based on local authority waste disposal activity reporting (www.wastedataflow.org) which provides an insight into the local shares of UK activity for recycling, landfilling, incineration and other treatment and disposal options. Waste incineration emissions are based on point source emissions data.

For some sources where regional data are not available, current local mapping grids have been used; these mapping grids are commonly based on census and other survey data that are periodically updated and used within UK emissions mapping and modelling work. For many sources, there is insufficient local data available back to 1990, and assumptions and extrapolations of available datasets have been used to present a time-series of air quality pollution emissions.

The inventories for England, Scotland, Wales and Northern Ireland aim to use the best available data. For most sources, more data are available in recent years than for 1990. For example, installation-specific fuel use

² The latest available data are taken from the December 2010 Energy Trends, <http://www.decc.gov.uk/en/content/cms/statistics/publications/trends/trends.aspx>

data from major industrial plant under EU ETS are available from 2005 onwards and data for sites regulated under IPC/IPPC are available from 1998 onwards for England and Wales, 1999 onwards for Northern Ireland, and in 2002 and from 2004 onwards for Scotland. These data sources are used where possible to inform back-casting of emission estimates. As such there remains a greater level of uncertainty in emission estimates from the earlier part of the time-series compared to more recent years. Furthermore, the data quality from these environmental regulatory systems has evolved over the years, as monitoring, reporting and quality checking methods and protocols have developed. This also impacts upon the accuracy of the reported emissions of AQ pollutants which are used within inventory compilation, such that more recent data are likely to be more accurate.

Air Quality Emission Inventories: Key Findings

The main findings of this report are summarised below:

ENGLAND

- Emissions of **ammonia** are estimated at 192kt in 2010. These emissions have declined by 22% since 1990 and account for 68% of the UK total in 2010. Agricultural sources dominate the inventory with manure management representing 64% of total ammonia emissions in 2010 and 35% coming from cattle manure management alone.
- Emissions of **carbon monoxide** are estimated at 1,645kt in 2010 and have declined by 77% since 1990. England's emissions account for 77% of the UK total. In 2010, 47% of emissions stem from road transport combustion sources.
- Emissions of **nitrogen oxides** are estimated at 826kt in 2010, representing 75% of the UK total in 2010. Emissions have declined by 64% since 1990, with 37% stemming from road transport combustion sources and 23% from power generation.
- Emissions of **non-methane volatile organic compounds** are estimated at 568kt in 2010, representing a 72% reduction in emissions since 1990. This reduction has been dominated by road transport, including evaporative losses, which have reduced by 93% since 1990. England represents 72% of the UK total.
- Emissions of **PM₁₀** are estimated at 86kt in 2010 and have declined by 59% since 1990. They account for 75% of the UK total. 24% of emissions come from road transport sources. Power generation accounted for 27% in 1990 but have been significantly reduced to 6% of England's total in 2010.
- Emissions of **sulphur dioxide** are estimated at 275kt in 2010, representing 68% of the UK total in 2010. Emissions have declined by 91% since 1990, which has been dominated by the 95% reduction in power generation due to the growth in gas and nuclear fuel use and the installation of FGD plant at a number of coal-fired power stations.
- Emissions of **lead** are estimated at 40t in 2010. Emissions have declined by 98% since 1990 and accounted for 68% of the UK total in 2010. The decline is dominated by the 1,799t reduction in transport sources due to the phase-out of leaded petrol. 30% of 2010 emissions arise due to the production in iron and steel industries.

SCOTLAND

- Emissions of **ammonia** are estimated at 35kt in 2010. These emissions have declined by 26% since 1990 and account for 12% of the UK total in 2010. Manure management represents 70% of total ammonia emissions in 2010, which has declined by 15% since 1990.
- Emissions of **carbon monoxide** are estimated at 189kt in 2010 and have declined by 77% since 1990. Scotland's emissions account for 9% of the UK total. Road transport combustion sources accounted for 33% of emissions in 2010. The sharp (17%) decrease in carbon monoxide emissions between 2008 and 2009 was predominantly driven by this sector.

- Emissions of **nitrogen oxides** are estimated at 110kt in 2010, representing 10% of the UK total in 2010. Emissions have declined by 61% since 1990, with 29% of total emissions in 2010 stemming from road transport combustion sources and a further 29% from power generation.
- Emissions of **non-methane volatile organic compounds** are estimated at 122kt in 2010, representing a 70% reduction in emissions since 1990. The food and drink sector represents 44% of emissions dominated by brewers and distilleries. Fugitive emissions from fuels make up 14% of the 2010 total and have reduced by 90% since 1990. Road transport emissions have been reduced by 94% since 1990.
- Emissions of **PM₁₀** are estimated at 13kt in 2010 and have declined by 57% since 1990. They account for 11% of the UK total. 24% of emissions come from commercial and residential combustion and emissions from power generation accounted for 25% in 1990 but have been reduced to 10% of Scotland's total in 2010.
- Emissions of **sulphur dioxide** are estimated at 81kt in 2010, representing 20% of the UK total in 2010. Emissions have declined by 73% since 1990, which has been dominated by the 70% reduction in power generation due to the growth in gas, renewable and nuclear fuel use. Recent fluctuating trends in electricity generation have had a noticeable impact on emissions.
- Emissions of **lead** are estimated at 3.8t in 2010. Emissions have declined by 98% since 1990 and accounted for 6% of the UK total in 2010. The decline is dominated by the >99% reduction in transport sources due to the phase-out of leaded petrol. 34% of 2010 emissions arise due to the energy industries.

WALES

- Emissions of **ammonia** are estimated at 27kt in 2010. These emissions have declined by 16% since 1990 and account for 9% of the UK total in 2010. Manure management represents 67% of total ammonia emissions in 2010, with 51% of total emissions originating from cattle manure management alone.
- Emissions of **carbon monoxide** are estimated at 190kt in 2010 and have declined by 72% since 1990. Wales' emissions account for 9% of the UK total. The iron & steel industry contributed to 33% of the 2010 total, decreasing by 65% since 1990. The additional industrial output from Wales during 2006 resulted in an increase in emissions between 2005 and 2006.
- Emissions of **nitrogen oxides** are estimated at 82kt in 2010, representing 7% of the UK total in 2010. Emissions have declined by 54% since 1990, with 29% of emissions in 2010 stemming from power generation. Recent trends in electricity generation have dominated the overall trends, with large fluctuations in coal-fired power generation.
- Emissions of **non-methane volatile organic compounds** are estimated at 43kt in 2010, representing a 69% reduction in emissions since 1990. This reduction has been dominated by the 93% decrease in road transport sources since 1990, as well as the 60% decline in the solvent and other product use sector. This sector accounted for 41% of emissions in 2010.
- Emissions of **PM₁₀** are estimated at 9kt in 2010 and have declined by 54% since 1990. They account for 8% of the UK total in 2010. 28% of emissions come from commercial and residential combustion and emissions from road transport sources accounted for 15%.
- Emissions of **sulphur dioxide** are estimated at 33kt in 2010, representing 8% of the UK total in 2010. Emissions have declined by 82% since 1990. Petroleum refineries are the most significant source in Wales, accounting for 37% of emissions. The installation of FGD at Aberthaw station has contributed to a reduction in emissions from power generation to only 12% in 2010, from 46% in 1990.
- Emissions of **lead** are estimated at 13.5t in 2010. Emissions have declined by 90% since 1990 and accounted for 23% of the UK total in 2010. The decline is dominated by the >99% reduction in transport sources due to the phase-out of leaded petrol. 81% of 2010 emissions came from industrial processes.

NORTHERN IRELAND

- Emissions of **ammonia** are estimated at 30kt in 2010. These emissions have declined by 8% since 1990 and account for 11% of the UK total in 2010. Manure management represents 79% of total ammonia emissions in 2010, with 61% of total emissions originating from cattle manure management alone.
- Emissions of **carbon monoxide** are estimated at 77kt in 2010 and have declined by 79% since 1990. Northern Ireland's emissions accounted for 4% of the UK total in 2010. Road transport combustion sources accounted for 34% of emissions in 2010, decreasing by 85% since 1990. Commercial, agricultural and residential combustion represented 48% of total emissions, showing a comparable reduction to road transport of 79% since 1990.
- Emissions of **nitrogen oxides** are estimated at 33kt in 2010, representing only 3% of the UK total in 2010. Emissions have declined by 66% since 1990, with 36% of total emissions in 2010 stemming from road transport. Industrial combustion accounted for 16% of 2010 emissions and 17% came from power generation.
- Emissions of **non-methane volatile organic compounds** are estimated at 19kt in 2010, representing a 71% reduction in emissions since 1990. This reduction has been dominated by the 94% decrease since 1990 in road transport sources, including evaporative losses. The solvent and other product use sector made up 51% of emissions in 2010.
- Emissions of **PM₁₀** are estimated at 5kt in 2010 and have declined by 67% since 1990. They account for 5% of the UK total in 2010. 37% of emissions came from commercial, residential and agricultural combustion, which has declined by 77% since 1990. Emissions from transport sources accounted for 19% of emissions in 2010.
- Emissions of **sulphur dioxide** are estimated at 14kt in 2010, representing 4% of the UK total in 2010. Power generation has dominated the 87% decline in sulphur dioxide emissions since 1990. Due to the sulphur in coal and fuel oil, power generation accounted for 60% of emissions in 1990 but has since been reduced by 97%. Residential combustion was the most significant source of emissions (51%) in 2010.
- Emissions of **lead** are estimated at 1.5t in 2010. Emissions have declined by 98% since 1990 and accounted for 3% of the UK total in 2010. The decline is dominated by the >99% reduction in transport sources due to the phase-out of leaded petrol. Domestic combustion accounted for 36% of emissions in 2010.

Per capita emissions

Emissions per capita have been calculated for each of the DAs, and are summarised for each pollutant within the report. Key features (e.g. where per capita emissions in one DA are much higher than the UK average) are described within the pollutant specific sections of the report, and a brief summary of the findings is described below:

- Across all DAs, for all pollutants, per capita emissions have fallen between 1990 and 2010;
- The most notable decrease (in percentage terms) is for lead, with a decrease of more than 90% across all of the DAs;
- In England, per capita emissions are lower than the UK average for all pollutants in 2010;
- In Northern Ireland, ammonia emissions per capita are almost four times the UK average in 2010. This is due to the very high contribution of emissions from agriculture, relative to the rest of the UK; Northern Ireland accounts for 11% of UK agriculture emissions, compared with just 3% of the UK population;
- Sulphur dioxide emissions per capita in Scotland are higher than the UK average, due to the high contribution of Scottish emissions from residential combustion and power stations to the UK totals for these sectors (21% and 34%, respectively, compared with only 8% of the UK population);
- Scottish VOC emissions per capita are 85% higher than the UK average, mostly due to the high contribution of Scottish emissions from food and drink manufacture to both the UK total for this sector, and the Scottish total VOC emission;

Air Quality Pollutants Inventories for England, Scotland
Wales and Northern Ireland: 1990-2010

- Welsh emissions are much higher than the UK average for NO_x , PM_{10} , SO_2 , and most notably for Pb and CO. This is due predominantly to the contribution of iron and steel industry emissions to the Welsh total.

Contacts

This work forms part of the Atmosphere & Local Environment (ALE) Programme of the Department for Environment, Food and Rural Affairs. AEA and Aether compile emission estimates for the energy, industrial process, solvents and waste sectors. The Centre for Ecology and Hydrology (Edinburgh) provides emission estimates for land use, land use change and forestry sources. Rothamsted Research provides the estimates of agricultural emissions.

Science policy enquiries should be directed to Sarah Honour, Atmosphere & Local Environment Programme, Department for Environment, Food and Rural Affairs, Ergon House, 17 Smith Square, London, SW1P 3JR, UK.

E-mail: agevidence@defra.gsi.gov.uk

Technical enquiries should be directed to Glen Thistlethwaite, AEA, The Gemini Building, Fermi Avenue, Didcot, Oxfordshire, OX11 0QR, UK.

E-mail: glen.thistlethwaite@aeat.co.uk

Technical enquiries on land use, land use change and forestry should be addressed to Amanda Thomson at the Centre for Ecology and Hydrology, Bush Estate, Penicuik, EH26 0QB, UK.

E-mail: amath@ceh.ac.uk

Technical enquiries on agriculture should be addressed to Tom Misselbrook, Rothamsted Research, Devon, EX20 2SB, UK.

E-mail: tom.misselbrook@rothamsted.ac.uk

A copy of this report and related data may be found on the Defra NAEI website:

<http://naei.defra.gov.uk/>

Contents

CONTACTS	VII
CONTENTS	VIII
1 INTRODUCTION	1
1.1 BACKGROUND TO INVENTORY DEVELOPMENT FOR THE DEVOLVED ADMINISTRATIONS	1
1.1.1 <i>Air quality emission reduction drivers</i>	1
1.2 INVENTORY METHODOLOGY & DATA AVAILABILITY	3
1.2.1 <i>Background: Data Availability and Inventory Uncertainty</i>	4
1.2.2 <i>Inventory Compilation Method</i>	5
1.2.2.1 NAEI Point Source Database.....	6
1.2.2.2 NAEI Emission Mapping Grids	7
1.2.2.3 Other Regional Data	7
1.3 REPORT STRUCTURE	9
2 AIR QUALITY POLLUTANTS	10
2.1 AMMONIA.....	10
2.1.1 <i>England Ammonia Inventory by Sector, 1990-2010</i>	12
2.1.2 <i>Scotland Ammonia Inventory by Sector, 1990-2010</i>	14
2.1.3 <i>Wales Ammonia Inventory by Sector, 1990-2010</i>	16
2.1.4 <i>Northern Ireland Ammonia Inventory by Sector, 1990-2010</i>	18
2.2 CARBON MONOXIDE	20
2.2.1 <i>England Carbon Monoxide Inventory by Sector, 1990-2010</i>	22
2.2.2 <i>Scotland Carbon Monoxide Inventory by Sector, 1990-2010</i>	25
2.2.3 <i>Wales Carbon Monoxide Inventory by Sector, 1990-2010</i>	28
2.2.4 <i>Northern Ireland Carbon Monoxide Inventory by Sector, 1990-2010</i>	31
2.3 NITROGEN OXIDES	33
2.3.1 <i>England Nitrogen Oxides Inventory by Sector, 1990-2010</i>	35
2.3.2 <i>Scotland Nitrogen Oxides Inventory by Sector, 1990-2010</i>	37
2.3.3 <i>Wales Nitrogen Oxides Inventory by Sector, 1990-2010</i>	40
2.3.4 <i>Northern Ireland Nitrogen Oxides Inventory by Sector, 1990-2010</i>	43
2.4 NON-METHANE VOLATILE ORGANIC COMPOUNDS	45
2.4.1 <i>England NMVOC Inventory by Sector, 1990-2010</i>	47
2.4.2 <i>Scotland NMVOC Inventory by Sector, 1990-2010</i>	50
2.4.3 <i>Wales NMVOC Inventory by Sector, 1990-2010</i>	53
2.4.4 <i>Northern Ireland NMVOC Inventory by Sector, 1990-2010</i>	56
2.5 PARTICULATE MATTER AS PM ₁₀	59
2.5.1 <i>UK Trends in PM₁₀ Emissions</i>	59
2.5.2 <i>England PM₁₀ Inventory by Sector, 1990-2010</i>	61
2.5.3 <i>Scotland PM₁₀ Inventory by Sector, 1990-2010</i>	63
2.5.4 <i>Wales PM₁₀ Inventory by Sector, 1990-2010</i>	65
2.5.5 <i>Northern Ireland PM₁₀ Inventory by Sector, 1990-2010</i>	68
2.6 SULPHUR DIOXIDE.....	70
2.6.1 <i>England Sulphur Dioxide Inventory by Sector, 1990-2010</i>	72
2.6.2 <i>Scotland Sulphur Dioxide Inventory by Sector, 1990-2010</i>	74
2.6.3 <i>Wales Sulphur Dioxide Inventory by Sector, 1990-2010</i>	77
2.6.4 <i>Northern Ireland Sulphur Dioxide Inventory by Sector, 1990-2010</i>	79
2.7 LEAD.....	81
2.7.1 <i>England Lead Inventory by Sector, 1990-2010</i>	83
2.7.2 <i>Scotland Lead Inventory by Sector, 1990-2010</i>	85
2.7.3 <i>Wales Lead Inventory by Sector, 1990-2010</i>	88

2.7.4	<i>Northern Ireland Lead Inventory by Sector, 1990-2010</i>	91
3	UNCERTAINTIES	93
3.1	AMMONIA.....	93
3.2	CARBON MONOXIDE.....	94
3.3	NITROGEN OXIDES.....	94
3.4	NON-METHANE VOLATILE ORGANIC COMPOUNDS.....	94
3.5	PM ₁₀	95
3.6	SULPHUR DIOXIDE.....	95
3.7	LEAD.....	95
4	REFERENCES	97

Appendices

Appendix A: Definition of NFR Codes and Sector Categories

Appendix B: Methods for Calculating Emission Distributions

List of Figures

Figure 2-1	UK Ammonia Emissions by Constituent Country, 1990-2010.....	10
Figure 2-2	England Ammonia Emissions by Sector, 1990-2010.....	12
Figure 2-3	Map of Ammonia Emissions in England, 2010.....	13
Figure 2-4	Scotland Ammonia Emissions by Sector, 1990-2010.....	14
Figure 2-5	Map of Ammonia Emissions in Scotland, 2010.....	15
Figure 2-6	Wales Ammonia Emissions by Sector, 1990-2010.....	16
Figure 2-7	Map of Ammonia Emissions in Wales, 2010.....	17
Figure 2-8	Northern Ireland Ammonia Emissions by Sector, 1990-2010.....	18
Figure 2-9	Map of Ammonia Emissions in Northern Ireland, 2010.....	19
Figure 2-10	UK Carbon Monoxide Emissions by Constituent Country, 1990-2010.....	20
Figure 2-11	England Carbon Monoxide Emissions by Sector, 1990-2010.....	22
Figure 2-12	Map of Carbon Monoxide Emissions in England, 2010.....	24
Figure 2-13	Scotland Carbon Monoxide Emissions by Sector, 1990-2010.....	25
Figure 2-14	Map of Carbon Monoxide Emissions in Scotland, 2010.....	27
Figure 2-15	Wales Carbon Monoxide Emissions by Sector, 1990-2010.....	28
Figure 2-16	Map of Carbon Monoxide Emissions in Wales, 2010.....	30
Figure 2-17	Northern Ireland Carbon Monoxide Emissions by Sector, 1990-2010.....	31
Figure 2-18	Map of Carbon Monoxide Emissions in Northern Ireland, 2010.....	32
Figure 2-19	UK Nitrogen Oxides Emissions by Constituent Country, 1990-2010.....	34
Figure 2-20	England Nitrogen Oxides Emissions by Sector, 1990-2010.....	35
Figure 2-21	Map of Nitrogen Oxides Emissions in England, 2010.....	36
Figure 2-22	Scotland Nitrogen Oxides Emissions by Sector, 1990-2010.....	37
Figure 2-23	Map of Nitrogen Oxides Emissions in Scotland, 2010.....	39
Figure 2-24	Wales Nitrogen Oxides Emissions by Sector, 1990-2010.....	40
Figure 2-25	Map of Nitrogen Oxides Emissions in Wales, 2010.....	42
Figure 2-26	Northern Ireland Nitrogen Oxides Emissions by Sector, 1990-2010.....	43
Figure 2-27	Map of Nitrogen Oxides Emissions in Northern Ireland, 2010.....	44
Figure 2-28	UK NMVOC Emissions by Constituent Country, 1990-2010.....	46
Figure 2-29	England NMVOC Emissions by Sector, 1990-2010.....	47
Figure 2-30	Map of NMVOC Emissions in England, 2010.....	49
Figure 2-31	Scotland NMVOC Emissions by Sector, 1990-2010.....	50
Figure 2-32	Map of NMVOC Emissions in Scotland, 2010.....	52

Figure 2-33	Wales NMVOC Emissions by Sector, 1990-2010.....	53
Figure 2-34	Map of NMVOC Emissions in Wales, 2010.....	55
Figure 2-35	Northern Ireland NMVOC Emissions by Sector, 1990-2010	56
Figure 2-36	Map of NMVOC Emissions in Northern Ireland, 2010	58
Figure 2-37	UK PM ₁₀ Emissions by Constituent Country, 1990-2010	60
Figure 2-38	England PM ₁₀ Emissions by Sector, 1990-2010.....	61
Figure 2-39	Map of PM ₁₀ Emissions in England, 2010	62
Figure 2-40	Scotland PM ₁₀ Emissions by Sector, 1990-2010.....	63
Figure 2-41	Map of PM ₁₀ Emissions in Scotland, 2010	64
Figure 2-42	Wales PM ₁₀ Emissions by Sector, 1990-2010.....	65
Figure 2-43	Map of PM ₁₀ Emissions in Wales, 2010	67
Figure 2-44	Northern Ireland PM ₁₀ Emissions by Sector, 1990-2010	68
Figure 2-45	Map of PM ₁₀ Emissions in Northern Ireland, 2010	69
Figure 2-46	UK Sulphur Dioxide Emissions by Constituent Country, 1990-2010	70
Figure 2-47	England Sulphur Dioxide Emissions by Sector, 1990-2010	72
Figure 2-48	Map of Sulphur Dioxide Emissions in England, 2010	73
Figure 2-49	Scotland Sulphur Dioxide Emissions by Sector, 1990-2010	74
Figure 2-50	Map of Sulphur Dioxide Emissions in Scotland, 2010	76
Figure 2-51	Wales Sulphur Dioxide Emissions by Sector, 1990-2010	77
Figure 2-52	Map of Sulphur Dioxide Emissions in Wales, 2010	78
Figure 2-53	Northern Ireland Sulphur Dioxide Emissions by Sector, 1990-2010.....	79
Figure 2-54	Map of Sulphur Dioxide Emissions in Northern Ireland, 2010	80
Figure 2-55	UK Lead Emissions by Constituent Country, 1990-2010.....	81
Figure 2-56	England Lead Emissions by Sector, 1990-2010	83
Figure 2-57	Map of Lead Emissions in England, 2010.....	84
Figure 2-58	Scotland Lead Emissions by Sector, 1990-2010.....	85
Figure 2-59	Map of Lead Emissions in Scotland, 2010.....	87
Figure 2-60	Wales Lead Emissions by Sector, 1990-2010	88
Figure 2-61	Map of Lead Emissions in Wales, 2010.....	90
Figure 2-62	Northern Ireland Lead Emissions by Sector, 1990-2010.....	91
Figure 2-63	Map of Lead Emissions in Northern Ireland, 2010.....	92

List of Tables

Table 2-1	Proportion of UK Ammonia Emissions by Constituent Country.....	11
Table 2-2	Emissions of Ammonia per capita by Constituent Country (kg/head)	11
Table 2-3	England Emissions of Ammonia by Sector, 1990-2010	12
Table 2-4	Scotland Emissions of Ammonia by Sector, 1990-2010	14
Table 2-5	Wales Emissions of Ammonia by Source, 1990-2010	16
Table 2-6	Northern Ireland Emissions of Ammonia by Sector, 1990-2010.....	18
Table 2-7	Proportion of UK Carbon Monoxide Emissions by Constituent Country.....	21
Table 2-8	Emissions of Carbon Monoxide per capita by Constituent Country (kg/head).....	21
Table 2-9	England Emissions of Carbon Monoxide by Sector, 1990-2010.....	22
Table 2-10	Scotland Emissions of Carbon Monoxide by Sector, 1990-2010.....	25
Table 2-11	Wales Emissions of Carbon Monoxide by Sector, 1990-2010	28
Table 2-12	Northern Ireland Carbon Monoxide Emissions by Sector, 1990-2010.....	31
Table 2-13	Proportion of UK Nitrogen Oxides Emissions by Constituent Country	34
Table 2-14	Emissions of NO _x per capita by Constituent Country (kg/head)	34
Table 2-15	England Emissions of Nitrogen Oxides by sector, 1990-2010	35
Table 2-16	Scotland Emissions of Nitrogen Oxides by sector, 1990-2010	37
Table 2-17	Wales Emissions of Nitrogen Oxides by sector, 1990-2010	40
Table 2-18	Northern Ireland Emissions of Nitrogen Oxides by sector, 1990-2010.....	43
Table 2-19	Proportion of UK NMVOC Emissions by Constituent Country.....	46
Table 2-20	Emissions of NMVOC per capita by Constituent Country (kg/head).....	46

Table 2-21	England Emissions of NMVOC by Sector, 1990-2010	47
Table 2-22	Scotland Emissions of NMVOC by Sector, 1990-2010	50
Table 2-23	Wales Emissions of NMVOC by Sector, 1990-2010	53
Table 2-24	Northern Ireland Emissions of NMVOC by Sector, 1990-2010	56
Table 2-25	Proportion of UK PM ₁₀ Emissions by Constituent Country	60
Table 2-26	Emissions of PM ₁₀ per capita by Constituent Country (kg/head)	60
Table 2-27	England Emissions of PM ₁₀ by Sector, 1990-2010	61
Table 2-28	Scotland Emissions of PM ₁₀ by Sector, 1990-2010	63
Table 2-29	Wales Emissions of PM ₁₀ by Sector, 1990-2010	65
Table 2-30	Northern Ireland Emissions of PM ₁₀ by Sector, 1990-2010	68
Table 2-31	Proportion of UK Sulphur Dioxide Emissions by Constituent Country	71
Table 2-32	Emissions of Sulphur Dioxide per capita by Constituent Country (kg/head)	71
Table 2-33	England Emissions of Sulphur Dioxide by Sector, 1990-2010	72
Table 2-34	Scotland Emissions of Sulphur Dioxide by Sector, 1990-2010	74
Table 2-35	Wales Emissions of Sulphur Dioxide by Sector, 1990-2010	77
Table 2-36	Northern Ireland Emissions of Sulphur Dioxide by Sector, 1990-2010	79
Table 2-37	Proportion of UK Lead Emissions by Constituent Country	82
Table 2-38	Emissions of Lead per capita by Constituent Country (g/head)	82
Table 2-39	England Emissions of Lead by Sector (tonnes), 1990-2010	83
Table 2-40	Scotland Emissions of Lead by Sector (tonnes), 1990-2010	85
Table 2-41	Wales Emissions of Lead by Sector (tonnes), 1990-2010	88
Table 2-42	Northern Ireland Emissions of Lead by Sector (tonnes), 1990-2010	91

1 Introduction

1.1 BACKGROUND TO INVENTORY DEVELOPMENT FOR THE DEVOLVED ADMINISTRATIONS

This study to develop AQ pollutant datasets for each of the constituent countries of the UK has been commissioned by Defra in order to better inform energy and environmental policy-makers within the Devolved Administrations in their pursuit of objectives set by the Air Quality Strategy for England, Scotland, Wales and Northern Ireland (AQS for ESWNI). These objectives also contribute to the UK's targets as a whole in terms of meeting both national and international targets on both local and transboundary air pollution.

The provision of DA-level datasets and subsequent identification of key sources at more regional and local levels is a key step to enable prioritisation of local action and to highlight the potential impacts of specific policies and measures. The time-series of AQ pollutant emissions provides an insight into the effects of environmental policies introduced since 1990, and may help to identify where win-win policies could be pursued to achieve both AQ and GHG policy goals.

1.1.1 Air quality emission reduction drivers

Overall air quality in the UK is currently estimated to be better than at any time since the industrial revolution. However air pollution is still estimated to reduce the life expectancy of every person in the UK by an average of 7-8 months (AQS, 2008). A number of policies are currently in place in the UK, which aim to improve air quality. This includes the national air quality strategy for England, Scotland, Wales and Northern Ireland.

Air Quality Strategy for England, Scotland, Wales and Northern Ireland

The original National Air Quality Strategy (NAQS) published in 1997 (DOE 1997) set out a framework of standards and objectives for the air pollutants of most concern (SO₂, PM₁₀, NO_x, CO, lead, benzene, 1, 3-butadiene and tropospheric ozone). The aim of the strategy was to reduce the air pollutant impact on human health by reducing airborne concentrations.

The NAQS identified air quality standards for eight priority pollutants based on the recommendations of the Expert Panel on Air Quality Standards (EPAQS, 1995) or World Health Organisation (WHO) guidance where no EPAQS recommendation existed. The NAQS has been subject to periodic review, with consultation documents being published in 1998 and 2001 (DETR 1998a, Defra 2001), and has subsequently evolved into the Air Quality Strategy for England, Scotland, Wales and Northern Ireland (AQS for ESWNI), with the same goals. A second edition of the strategy was published in 2000 (DETR 2000), identifying further revisions and focused on the incorporation of air quality limit values in European Directives, and the impacts of devolution. On 17 July 2007 a new Air Quality Strategy was published by Defra and the Devolved Administrations. The details of this AQS can be found on the Defra website at: <http://www.defra.gov.uk/publications/tag/air-quality-strategy/>.

The new Air Quality Strategy supersedes previous versions and covers the whole of the UK, therefore including DA-specific objective values that were previously detailed in addenda to the previous AQS.

EU Air Quality Framework Directive

The EU air quality framework directive (96/62/EC) established a framework for setting limit values, assessing concentrations and managing air quality to avoid exceeding the limits for air pollutants known to be harmful to human health and the environment through a series of four Daughter Directives. However, in 2008, the Framework Directive and first three Daughter Directives were consolidated in a new EU air quality Directive (2008/50/EC), the 4th Daughter Directive. The new Directive also introduced a new regulatory framework for PM_{2.5}.

At present, under 2008/50/EC and the 4th Daughter Directive, limit values are set for twelve pollutants, including NO_x, SO₂, PM and CO, and member states are required to submit annual reports to the European Commission on whether the limits have been achieved within their respective areas.

UNECE Convention on Long-Range Transboundary Air Pollution

The UK is committed to reducing acidifying gas and ozone precursor emissions and is a party to several protocols under the UNECE Convention on Long-Range Transboundary Air Pollution.

Under the Second Sulphur Protocol, the UK committed to reducing its total SO₂ emissions by 50% by 2000, 70% by 2005 and 80% by 2010 (all from a 1980 baseline).

The NMVOC Protocol requires the UK to achieve a 30% reduction of anthropogenic NMVOC emissions by 1999 from a 1988 baseline. The emission estimates given in the 1999 version of the emissions inventory indicated that this was achieved.

The NO_x Protocol required that the total emissions of NO_x in 1994 should be no higher than they were in 1987; UK emissions were 11% lower in 1994 than in 1987 and have fallen substantially since 1994.

In 1996, the UNECE started negotiating a new multi-effect, multi-pollutant protocol on nitrogen oxides and related substances. This was aimed at addressing photochemical pollution, acidification and eutrophication. The Protocol to Abate Acidification, Eutrophication and Ground-level Ozone was adopted in Gothenburg in December 2000, where it was signed by the UK. The multi-pollutant protocol incorporates several measures to facilitate the reduction of emissions:

- Emission ceilings are specified for sulphur, nitrogen oxides, NH₃ and NMVOCs;
- Emission limits are specified for sulphur, nitrogen oxides and NMVOCs from stationary sources;
- Emission limits are indicated for CO, hydrocarbons, nitrogen oxides and particulates from new mobile sources;
- Environmental specifications for petrol and diesel fuels are given;
- Several measures to reduce NH₃ emissions from the agriculture sector (4) are required.

The Gothenburg Protocol forms a part of the Convention on Long-range Transboundary Air Pollution. More detailed information on both of the Gothenburg protocol and the Convention may be found at the UNECE web site: www.unece.org/env/lrtap/

National Emissions Ceilings Directive

Within the EU, the National Emission Ceilings Directive was agreed in 2001. It sets emission ceilings to be achieved from 2010 onwards for each Member State for the same four pollutants as in the Gothenburg Protocol. The UK has met current international targets to reduce total emissions by 2010 of four air pollutants that cause harm to people's health and to the natural environment:

- Emissions of **sulphur dioxide** have fallen by 89 per cent between 1990 and 2010, but increased by 2 per cent between 2009 and 2010. Emissions in 2010 were 31 per cent below the lowest international target for the UK.
- Emissions of **nitrogen oxides** have fallen by 62 per cent between 1990 and 2010, and fell by 3 per cent between 2009 and 2010. Emissions in 2010 were 5 per cent below the lowest international target for the UK.
- Emissions of **non-methane volatile organic compounds** have fallen by 71 per cent between 1990 and 2010, and fell by 4 per cent between 2009 and 2010. Emissions in 2010 were 34 per cent below the international target for the UK.
- Emissions of **ammonia** have fallen by 21 per cent between 1990 and 2010, but increased by 0.5 per cent between 2009 and 2010. Emissions in 2010 were 4 per cent below the international target for the UK.

More information on these ceilings can be found in a statistical release from Defra (2011).

Large Combustion Plant Directive

Within the UK, the implementation of the EC's Large Combustion Plant Directive and other associated policy measures has led to substantial reductions in acidifying pollutants, specifically NO_x, SO₂ and dust from power plants and industrial sources.

Sulphur Content of Liquid Fuels Directive

The EC's Directive to limit sulphur content in gas oil and fuel oil has been transposed into UK regulations which were initially established in 2000 but were updated with Statutory Instruments brought into force across the DAs via the Sulphur Content of Liquid Fuel Regulations 2007 (England and Wales: SI79/2007; Scotland: SI 27/2007; Northern Ireland: SI 272/2007). The main impact of these regulations has been to gradually drive down the maximum sulphur content of refinery products, with the 2007 Regulations requiring that gas oil has a maximum 0.1% content Sulphur by mass from January 2008 onwards. The impacts of this change are evident within the recent emission trends of the UK and DA inventories as SO₂ emissions have declined significantly between 2007 and 2008 from road transport (1A3b) and other sources where petroleum-based fuels are dominant.

UNECE Heavy Metals Protocol

The Convention on Long-range Transboundary Air Pollution was signed in 1979 and came into force in 1983. Since its entry into force, the Convention has been extended by a number of protocols, including the 1998 Protocol on Heavy Metals. This Protocol is given in outline below; more information may be found at the UN/ECE web site, located at: <http://www.unece.org/env/lrtap/>. The UK has signed this protocol.

The UNECE Protocol on Heavy Metals targets three particularly harmful substances: lead, cadmium and mercury. Countries are obliged to reduce their emissions of these three metals below their levels in 1990 (or an alternative year between 1985 and 1995). The protocol aims to cut emissions from industrial sources (iron and steel industry, non-ferrous metal industry), combustion processes (power generation, road transport) and waste incineration.

The protocol specifies limit values for emissions from stationary sources and requires the use of Best Available Technology (BAT) to minimise emissions from these sources, through the application of special filters or scrubbers for combustion sources, or mercury-free processes. The protocol also required countries to phase out leaded petrol. Under the protocol, measures are introduced to lower heavy metal emissions from other products e.g. mercury in batteries, and examples are given of management measures for other mercury containing products, such as electrical components (thermostats, switches), measuring devices (thermometers, manometers, barometers), fluorescent lamps, dental amalgam, pesticides and paint.

1.2 INVENTORY METHODOLOGY & DATA AVAILABILITY

This report presents emission inventories for the constituent countries of the UK for the period 1990 to 2010, for the following priority Air Quality (AQ) pollutants:

- Ammonia (NH₃)
- Carbon monoxide (CO)
- Nitrogen oxides (reported as nitrogen dioxide) (NO_x as NO₂)
- Non-methane volatile organic compounds (NMVOC)
- Sub-10 micron particulate matter (PM₁₀)
- Sulphur dioxide (SO₂)
- Lead (Pb)

The estimates have been compiled by disaggregating the UK emission totals presented within "UK Informative Inventory Report 1980 to 2010" (Passant *et al.*, 2011), derived from the National Atmospheric Emissions Inventory database. The UK data is compiled annually in accordance with the requirements of United Nations Economic Commission for Europe (UNECE) reporting guidelines using the NFR reporting format and submitted to the Convention on Long-Range Transboundary Air Pollution (CLRTAP).

1.2.1 Background: Data Availability and Inventory Uncertainty

The method for disaggregating UK emission totals across the constituent countries draws on a combination of point source data (e.g. Pollution Inventory³ data for industrial emissions) and sub-national and local datasets such as:

- DECC sub-national statistics on energy use;
- Other regional energy use data for specific industries or regional data on raw material consumption or sector-specific production;
- Major road traffic count data;
- Domestic and international flight data for all major UK airports;
- Rail company fuel use estimates;
- Regional housing, employment, population and consumption data;
- Agricultural surveys (livestock numbers, crop production, fertiliser application);
- Land use survey data.

Emissions from the offshore oil & gas exploration and production sector are not attributed to a specific country inventory, but are reported within an “unallocated” category. Note, however, that emissions from onshore oil & gas terminals are assigned to the appropriate country inventories.

The disaggregation of air quality (AQ) pollutant emissions across the four constituent countries of the UK has been conducted five times previously and this report presents the results from a programme of on-going data and methodology improvement, to provide emission inventories for the Devolved Administrations (DAs). This programme spans both GHG and AQ emission inventories, and is driven by the developing requirements for sub-national reporting against emission targets and DA policy development.

For many emission sources of AQ pollutants, the data available for constituent country emissions are less detailed than for the UK as a whole, and for some sources country-level data are not available at all. For this reason, a “top-down” approach using UK inventory data as the core dataset has been adopted, and percentage splits of the UK total have been derived for each of the constituent countries using available regional data.

In particular, energy balance data (i.e. fuel production, transformation and sector-specific consumption data) are not available for England, Wales and Scotland. Sub-national energy statistics are published annually by the Department for Energy and Climate Change (DECC) within the quarterly Energy Trends⁴ publication. These sub-national statistics are limited in their detail when compared to UK-level energy statistics, but do provide estimated fuel use data for England, Scotland, Wales and Northern Ireland for the following source sectors:

- Industry (1A2) & Commercial (1A4a)
- Agriculture (combustion source) (1A4c)
- Residential (1A4b)

The DECC sub-national energy statistics have been developed in recent years to provide estimates of fuel use and CO₂ emissions data at Local Authority (LA) level across the UK. The latest available data include LA solid and liquid fuel use estimates for 2005 to 2009, with gas and electricity data also being available up to 2010.

The DECC data at local and regional level are derived from analysis of gas and electricity meter point data, supplemented by additional research to estimate the distribution of solid fuels and petroleum-based fuels across the UK. Since the initial study and presentation of experimental data for 2003 and 2004, each annual revision to the local and regional data has included data improvements through targeted sector research. These DECC sub-national energy statistics continue to evolve and improve, reducing data inaccuracies, and are the best data available to inform the patterns of fuel use across the Devolved Administrations. They are

³ The term “Pollution Inventory” is used here to represent the industrial emissions databases of the UK environmental regulators (The Environment Agency of England & Wales, the Scottish Environment Protection Agency and the Northern Ireland Department of Environment) which comprise annual emission estimates from all IPC/IPPC-regulated processes under their authority.

⁴ The latest available data are taken from the December 2011 Energy Trends, <http://www.decc.gov.uk/en/content/cms/statistics/publications/trends/trends.aspx>

therefore used to underpin the pollutant emission estimates from fuel combustion sources within the inventories presented here, in conjunction with other data sources such as EU ETS fuel use data for large industrial sites and other DA-specific energy data.

For other significant emission sources there are complete country-level datasets available, although some of these are less detailed than data used for the UK Inventory:

- Industrial process emissions are based on plant operator estimates reported to environmental agencies under regulatory systems such as Integrated Pollution Prevention and Control (IPPC). Major sources include power stations, cement and lime kilns, iron & steelworks, aluminium and other non-ferrous metal plant, chemical industries. These data are not available across the full time series from 1990, as the regulatory reporting regimes developed in the late 1990s (in England, Wales and Northern Ireland) and early 2000s (in Scotland);
- Emissions from oil and gas terminals and offshore platforms and rigs, are based on operator estimates reported to the DECC Oil and Gas team (2011) in Aberdeen through the Environmental Emissions Monitoring System, EEMS;
- Agricultural emissions are based on UK emission factors and annual survey data across each of the Devolved Administrations, including estimates of arable production and livestock numbers (Rothamsted Research, 2011);
- Emissions from waste disposal activities are estimated based on modelled emissions from the UK air quality inventory (Defra, 2012) split out across the DAs based on local authority waste disposal activity reporting (www.wastedataflow.org) which provides an insight into the local shares of UK activity for recycling, landfilling, incineration and other treatment and disposal options. Waste incineration emissions are based on point source emissions data.
- For some sources where regional data are not available, current local mapping grids have been used; these mapping grids are commonly based on census and other survey data that are periodically updated and used within UK emissions mapping and modelling work (Tzagatakis *et al.*, 2011).

In many source sectors, there is insufficient local data available back to 1990 or earlier, and assumptions and extrapolations of available datasets have frequently been used to present a time-series of air quality pollution emissions.

As a result of the more limited country-specific activity and emission factor data, the emission estimates for the England, Scotland, Wales and Northern Ireland inventories are subject to greater uncertainty than the equivalent UK estimates. There are step-changes in data availability during the time-series, such as installation-specific fuel use data from major industrial plant under EU ETS (from 2005 onwards) and sites regulated under IPC/IPPC (1998 onwards for England and Wales, 1999 onwards for Northern Ireland, and in 2002 and from 2004 onwards for Scotland). These data sources are used where possible to inform back-casting of emission estimates, but there remains a greater level of uncertainty in emission estimates from the earlier part of the time-series compared to more recent years. Furthermore, the data quality from these environmental regulatory systems has evolved over the years, as monitoring, reporting and quality checking methods and protocols have developed. This also impacts upon the accuracy of the reported emissions of AQ pollutants which are used within inventory compilation, such that more recent data are likely to be more accurate. The uncertainties in the DA air quality inventories are discussed in more detail in Chapter 3.

1.2.2 Inventory Compilation Method

A comprehensive list of all sources and UK emissions for the target pollutants (CO, NO_x, SO₂, VOC, NH₃, PM₁₀, Pb) during the study period of 1990-2010 is available from the NAEI database. From these data, the key sources for each of the AQ pollutants can be determined. The DA share of the UK emissions from each source category are then determined using the best available regional data, which may range from good quality emissions or activity data, to the use of proxy data (e.g. production or employment indices, population data) to provide a “best estimate” of the DA share of the UK emissions from a given source.

There are a number of resources that have been used to analyse the DA share of UK emissions for each emission source, including:

- NAEI point source database;
- Emissions mapping grid data;
- Regional data derived from analysis of activity data trends, taken from research to develop DA Greenhouse Gas (GHG) Inventories;
- Generic parameters and proxy data such as population or regional GDP data.

The development of more consistent reports and datasets between different scales (national-regional-local) derived from the NAEI database is a key improvement that this study has enabled. The main resources used within the DA air quality pollutant inventory analysis are outlined below.

1.2.2.1 NAEI Point Source Database

Operators of all IPC/IPPC-regulated industrial plant are required to submit annual emission estimates of a range of pollutants (including all of those pertinent to this study) to their local UK environmental regulatory agency, and these emission estimates are subject to established procedures of Quality Assurance and Quality Checking prior to publication. These industrial point-source pollution inventories (held by the Environment Agency, the Scottish Environment Protection Agency and the Northern Ireland Environment Agency) are emission datasets that have been developing and improving since their inception in the mid-1990s. Robust and reliable data for installations in England and Wales have been widely available since around 1998, whilst the equivalent datasets in Scotland and Northern Ireland became available from the early 2000s.

NAEI point source data have been improved over recent years through the increasing quality and availability of these IPC/IPPC-regulated industrial pollution emission datasets, as well as through the availability of site-specific fuel use data for sites that operate within the EU Emissions Trading System (EU ETS), which has been running since 2005. Annual data requests are also made directly to plant operators in key sectors such as power stations, refineries, cement & lime manufacture, iron & steel manufacture, chemical industry and waste treatment and disposal, in order to procure more detailed emissions data and other parameters (e.g. production data).

As part of the Devolved Administrations Inventory Improvement Programme, a research study was undertaken in early 2010 to source more detailed information on emissions sources at a number of petrochemical and industrial sites across the UK. The study included consultation with Environment Agencies responsible for each of the respective Devolved Administrations and site visits to review further details of applications and reports submitted in relation to permitted activities.

Through analysis of the time-series of data and review of the latest emission estimates, the point source data is amended as appropriate to fill in gaps and rectify any errors. These finalised data are then used as the basis for the NAEI industrial emissions estimates. The location of each site is known and therefore the point-source database can be queried to extract all emissions information relevant to a given geographical area, and hence the DA-level inventories can partly be populated in this way.

Although the use of this dataset can only provide a limited time-series of emissions from a given source sector, it is nevertheless a useful tool for deriving recent regional emissions data for a broad range of pollutants, including CO, NO_x, SO₂, VOC, Pb, NH₃ and PM₁₀. The NAEI point-source database is most useful for industries that are dominated by large IPC/IPPC-authorized plant, such as power stations, refineries, iron & steel manufacturing, cement and lime kilns and so on. For these sectors, the point source database covers nearly 100% of emissions, and is regarded to be the best available dataset for such sources, as it is largely based on energy use and emissions data derived from regulatory agency sources that are subject to quality checking and (in the case of EU ETS data) independent verification.

Annual revisions to the NAEI point source database are conducted when new data become available and/or when installation-level data are revised by operators, regulators or through enquiry by the AEA inventory team to resolve data discrepancies which may be evident between reporting mechanisms.

1.2.2.2 NAEI Emission Mapping Grids

Emission maps for the whole of the UK are routinely produced as part of the NAEI for 25 pollutants, including all of the pollutants considered in this study. The maps are compiled at a 1km resolution and are produced annually for the most recent NAEI database (2010 in this case). The mapped emissions data are available on the NAEI web site at:

http://naei.defra.gov.uk/mapping/mapping_2010.php

The emission maps are used by AEA and other organisations for a variety of Government policy support work at the national scale. In particular, the maps are used as input into a programme of air pollution modelling studies.

The geographical distribution of emissions across the UK is built up from distributions of emissions in each source sector. These source sector distributions are developed using a set of statistics appropriate to that sector. For large industrial 'point' sources, emissions are compiled from a variety of official UK sources (Environment Agency, Scottish Environment Protection Agency, Northern Ireland Environment Agency Local Authority data). For sources that are distributed widely across the UK (known as 'area' sources), a distribution map is generated using appropriate surrogate statistics for that sector. The method used for each source varies according to the data available, but is commonly based on either local activity statistics such as raw material use, energy use, industrial production and employment data, housing and population data, road vehicle and fuel sales data, periodic census or socio-economic survey data.

Periodic surveys and censuses of industrial, commercial, domestic, and other economic sectors provide indicators regarding the location and scale of a wide variety of activity data that can be used to disaggregate emissions totals, and these are commonly utilised within the NAEI mapping grids. For a more detailed description of the integration of point source data analysis and the development of UK emission maps, see Chapter 3 of *UK Emission Mapping Methodology 2009* (Tsagatakis et al., 2011). Appendix B of this report provides a summary table of the mapping grid data availability for each UNECE sector. Changes to the mapping grids used in the 1990-2010 inventory include:

- An updated distribution grid for adhesive manufacture;
- A move away from using a mapping grid for aluminium production and slag cement production to use a time series of data (derived from point source data);
- A change to using the mapping grid for spirits to reflect malt production (instead of malt exports);
- A change to using point source data for secondary aluminium production.

The key limitation to the use of mapping grids within inventory development is the difficulty in obtaining an accurate time-series of emissions from a given sector, as the mapping grids are typically only updated every few years as more survey data becomes available. The data availability limitations inevitably impact upon the reliability of emission inventory estimates. In this study we have endeavoured to focus resources on ensuring that the most significant sources are assessed most accurately across the time series, whilst less significant source sectors may be disaggregated using a mapping grid for all years in the time series.

1.2.2.3 Other Regional Data

In recent years, the NAEI team has aimed to develop a consistent time-series of detailed datasets to inform DA and local emission inventories (back to 1990) and pollutant mapping campaigns. Examples of such datasets that have been used in this study include:

- Sub-national fuel use data for natural gas, solid fuel and petroleum-based fuels, from UK Transco (Transco, 2011), other gas network operators, the Coal Authority (Coal Authority, 2011) and the Department of Energy and Climate Change (DECC). The AEA energy mapping team has been involved in the on-going development of the DECC sub-national energy statistics which provide limited data from 2004 to 2009. These data are used to underpin many of the AQ pollutant emission estimates from small-scale (non-regulated) combustion sources such as domestic, commercial, public administration and small-scale industrial sectors. Back-casting the fuel use trends to 1990 has drawn upon available UK-level data and trends supplemented by analysis of additional data, such as Housing

Condition Survey data, to ensure that significant changes are represented in the inventories (e.g. to reflect the development of the gas supply infrastructure in Northern Ireland since 1999). A reallocation of gas oil was applied to the inventory leading to improved consumption estimates for industrial combustion (1A2) and oil and gas extraction (1A1).

- The Road Transport (1A3b) emissions database uses local traffic count data from the Department for Transport (DfT), the Northern Ireland Department of Regional Development (DRDNI), fuel use datasets (DECC), vehicle fleet data (DfT, DRDNI) and emission factors developed by TRL on behalf of DfT and from European research sources (COPERT III, IV) (EEA, 2010b) to derive detailed emission estimates for a wide range of pollutants across the UK. There were a number of improvements made to the 2010 UK road transport inventory and thus affecting the DA inventories including the application of Automatic Number Plate Recognition (ANPR) data and revised emission factors for NO_x for all vehicle types (except motorcycles). These and other changes are described in detail in Appendix B.
- Aircraft emissions are derived from the Civil Aviation Authority's (CAA, 2011) database of flight movements, fuel use data (DECC), aircraft fleet information (CAA, 2011) and emission factors from international guidance and research (Intergovernmental Panel on Climate Change, IPCC) to derive emission estimates for aircraft cruise, take-off and landing cycles.
- Regional quarry production data and quarry location information (British Geological Survey) (BGS, 2011).
- Regional iron & steel production data, and regional fuel use data in the iron & steel industry (Tata Steel, 2011), (ISSB, 2011).
- Site-specific emissions data split by combustion and process sources for all UK refineries, and refinery production capacities (UKPIA, 2011).
- Site-specific cement production capacities and UK-wide cement industry fuel use data (British Cement Association, 2011).
- Regional railway diesel consumption data (local train operating companies, including freight, intercity and local passenger services).
- Regional housing & population data (Department of Communities and Local Government).
- Regional economic activity & industrial production indices (Office of National Statistics) (ONS, 2011).

1.3 REPORT STRUCTURE

This report is structured as follows:

Main body of the report: This part of the report presents and discusses the inventories for England, Scotland, Wales and Northern Ireland, providing air quality pollutant emissions data for the years 1990, 1995, and 1998 to 2010. Emission inventories for PM₁₀, CO, NMVOCs, NH₃, NO_x, SO₂ and Pb are included in Chapter 2. Where appropriate, the reasons for any significant trends in emissions, issues regarding data availability and uncertainty estimates are provided for each inventory. A qualitative assessment of the uncertainty in the DA air quality inventories is presented in Chapter 3.

Appendix A: This appendix provides National Reporting Format (NFR) sector code descriptions.

Appendix B: This appendix provides a summary of the disaggregation methods and mapping grids used in this study, for each UNECE sector.

2 Air Quality Pollutants


Inventories for England, Scotland, Wales and Northern Ireland for ammonia (NH₃), carbon monoxide (CO), nitrogen oxides (NO_x), non-methane volatile organic compounds (NMVOCs), sub-10 micron particulate matter (PM₁₀), sulphur dioxide (SO₂) and lead (Pb) are discussed in the following sections. These data have been derived by disaggregation of the UK figures using point source, mapping and regional datasets as appropriate (see Appendix B for details).

2.1 AMMONIA

Ammonia (NH₃) emissions play an important role in a number of different environmental issues including acidification, eutrophication and changes in biodiversity. The atmospheric chemistry of NH₃ and NH₄⁺ is such that transport of the pollutants can vary greatly, and that as a result, NH₃ emissions can exert impacts on a highly localised level, as well as contributing to the effects of long-range pollutant transport.

UK emission estimates for NH₃ are only available from 1990 onwards, because earlier data from the most significant industrial sources are not available for use in emission inventory estimates. UK ammonia emissions in 2010 represent a decrease of 21% on the 1990 emissions (**Figure 2-1**).

Figure 2-1 UK Ammonia Emissions by Constituent Country, 1990-2010



The main source of NH₃ emissions in the UK is livestock manure management, and in particular cattle manure management. These emissions derive mainly from the decomposition of urea in animal wastes and uric acid in poultry wastes. Emissions from nitrogen fertiliser use on grassland and arable crops are also a significant source and included in the ammonia inventory. Emissions are affected by a large number of factors, including animal species, age, weight, diet, housing and manure management systems, and environmental conditions. As such, the interpretation and extrapolation of experimental data is problematic, making emission estimates uncertain. Estimates are based on official livestock datasets, annual fertiliser use surveys, farm management practice surveys and detailed emission factors from recent literature sources. The methodology for compiling

the inventory of ammonia emissions from agriculture follows that of Webb and Misselbrook (2004) for manure management sources and Misselbrook et al (2004) for fertiliser sources, with annual revisions to input data, emission factors and other parameters as described in the annual Informative Inventory Report.

Decreasing livestock numbers (cattle in particular) and fertiliser use in the UK since 1990 have led to reductions in UK ammonia emissions, and it is this trend in agricultural sources that influences the DA-level inventories most significantly.

Non-agricultural sources of ammonia comprise a number of diverse sources. Emission estimates for these sources are often highly uncertain due to a lack of activity and emission factor data. Emissions from road transport (1A3b) (although relatively insignificant compared to agricultural emissions) increased in the 1990's as a result of the increasing number of three way catalysts in the vehicle fleet. However, emissions are now falling as the second generation of catalysts (which lead to lower NH₃ emissions than first generation catalysts) penetrates the vehicle fleet.

Emissions of ammonia for England, Wales, Scotland and Northern Ireland are summarised in the tables and graphs below. **Table 2-1** shows how the estimated total UK NH₃ emissions are split between the four constituent countries, and **Table 2-2** show emissions of ammonia per capita, with a comparison against the average for the UK.

Table 2-1 Proportion of UK Ammonia Emissions by Constituent Country

Year	England	Scotland	Wales	Northern Ireland	Unallocated
1990	69%	13%	9%	9%	0%
2010	68%	12%	9%	11%	0%

Table 2-2 Emissions of Ammonia per capita by Constituent Country (kg/head)

Year	England	Scotland	Wales	Northern Ireland	UK
1990	5.2	9.4	11.1	20.7	6.3
2010	3.7	6.7	8.9	16.9	4.6

2.1.1 England Ammonia Inventory by Sector, 1990-2010

The table and graph below give a summary of the ammonia emissions in England by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-3 England Emissions of Ammonia by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Transport Sources	0.7	5.8	11.5	13.3	19.3	18.0	16.9	15.4	14.4	13.3	12.3	11.3	10.1	9.6	8.6
Commercial, domestic and agricultural combustion	3.1	2.2	2.0	2.1	1.7	1.7	1.3	1.2	1.1	1.0	0.9	1.0	1.1	1.1	1.2
Industrial Processes	7.1	7.2	8.8	4.8	3.5	3.6	3.5	3.2	3.1	5.3	5.0	4.8	4.2	4.0	3.9
Agriculture	227.3	209.0	201.1	202.1	189.7	186.6	183.5	175.8	182.1	177.8	178.9	174.1	165.3	166.1	167.3
Other ¹	9.2	10.8	10.5	10.7	10.8	11.0	10.9	10.7	10.5	10.6	10.8	10.9	11.3	11.6	11.3
Total	247.4	234.9	234.0	233.0	225.0	220.8	216.1	206.3	211.3	207.9	208.0	202.2	192.0	192.4	192.2

Units: kilotonnes

¹ "Other" for ammonia includes emissions from Energy Industries, Industrial Combustion, Fugitive, Solvent Processes, Waste as well as 1A5b, 7A.


Figure 2-2 England Ammonia Emissions by Sector, 1990-2010



England's ammonia emissions have declined by 22% since 1990 and currently account for 68% of the UK total. The inventory is dominated by emissions from agricultural sources with 64% of the total in 2010 coming from manure management (4B: down 25% since 1990). 35% of the English total is from cattle manure management alone (4B1: down 19% since 1990). Other sources of note include transport emissions (1A3: 4% of the England total in 2010) and waste treatment and disposal (6: 5% of the England total in 2010).

English ammonia emissions per capita are much lower than the UK average. This is mostly due to the smaller contribution of emissions from agriculture in England; English emissions from this sector account for 66% of UK emissions for the sector, which is much lower than England's contribution to the population (84%).

Figure 2-3 Map of Ammonia Emissions in England, 2010



2.1.2 Scotland Ammonia Inventory by Sector, 1990-2010

The table and graph below give a summary of the ammonia emissions in Scotland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-4 Scotland Emissions of Ammonia by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Transport Sources	0.07	0.60	1.13	1.29	1.92	1.79	1.69	1.55	1.44	1.32	1.24	1.13	1.00	0.94	0.84
Commercial, domestic and agricultural combustion	0.80	0.45	0.40	0.42	0.35	0.33	0.29	0.25	0.23	0.19	0.24	0.27	0.29	0.30	0.32
Industrial Processes	0.08	0.08	0.08	0.07	0.07	0.07	0.06	0.06	0.07	0.08	0.06	0.07	0.06	0.06	0.07
Agriculture	45.55	41.12	38.50	38.95	36.70	37.05	36.36	35.06	35.99	35.35	34.72	33.03	31.75	31.53	32.18
Other ¹	1.29	1.52	1.48	1.50	1.51	1.53	1.52	1.48	1.44	1.48	1.50	1.57	1.65	1.82	1.80
Total	47.79	43.77	41.59	42.24	40.54	40.77	39.92	38.40	39.17	38.42	37.77	36.07	34.74	34.65	35.20

Units: kilotonnes


¹ "Other" for ammonia includes emissions from Energy Industries, Industrial Combustion, Fugitive, Solvent Processes, Waste as well as 1A5b, 7A.

Figure 2-4 Scotland Ammonia Emissions by Sector, 1990-2010



Scotland's ammonia emissions have declined by 26% since 1990 and accounted for 12% of the UK total in 2010. The inventory is dominated by emissions from agricultural sources, with 70% of the total in 2010 estimated to originate from manure management (4B: down 15% since 1990). Other sources of note include transport emissions (1A3: 2% of the Scotland total in 2010) and waste treatment and disposal (6: 4% of the Scotland total in 2010).

Figure 2-5 Map of Ammonia Emissions in Scotland, 2010



2.1.3 Wales Ammonia Inventory by Sector, 1990-2010

The table and graph below give a summary of the ammonia emissions in Wales by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-5 Wales Emissions of Ammonia by Source, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Transport Sources	0.04	0.39	0.72	0.82	1.22	1.13	1.07	0.98	0.92	0.84	0.79	0.72	0.64	0.60	0.53
Commercial, domestic and agricultural combustion	0.57	0.45	0.41	0.44	0.35	0.35	0.27	0.24	0.24	0.21	0.23	0.26	0.28	0.29	0.31
Industrial Processes	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.05	0.05	0.07	0.07	0.06	0.05	0.06
Agriculture	30.26	28.51	27.82	28.52	27.31	27.68	26.15	27.35	27.33	27.26	27.85	25.44	23.38	23.50	24.48
Other ¹	0.75	0.84	0.83	0.86	0.85	0.92	0.98	0.94	0.87	0.89	0.95	1.00	1.10	1.24	1.25
Total	31.68	30.25	29.84	30.69	29.78	30.13	28.52	29.55	29.40	29.25	29.88	27.49	25.47	25.68	26.63

Units: kilotonnes


¹ "Other" for ammonia includes emissions from Energy Industries, Industrial Combustion, Fugitive, Solvent Processes, Waste as well as 1A5b, 7A.

Figure 2-6 Wales Ammonia Emissions by Sector, 1990-2010



Ammonia emissions in Wales have declined by 16% since 1990 and accounted for 9% of the UK total in 2010. The inventory is dominated by emissions from agricultural sources with 67% of the total in 2010 coming from manure management (4B: down 13% since 1990). In 2010, 51% of emissions from Wales are estimated to originate from cattle manure management alone (4B1: down 12% since 1990). Other sources of note include transport emissions (1A3: 2% of the Wales total in 2010) and waste treatment and disposal (6: 4% of the Wales total in 2010).

Figure 2-7 Map of Ammonia Emissions in Wales, 2010



2.1.4 Northern Ireland Ammonia Inventory by Sector, 1990-2010

The table and graph below give a summary of the ammonia emissions in Northern Ireland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-6 Northern Ireland Emissions of Ammonia by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Transport Sources	0.03	0.27	0.49	0.56	0.88	0.80	0.72	0.65	0.62	0.57	0.52	0.49	0.44	0.43	0.38
Commercial, domestic and agricultural combustion	0.62	0.38	0.31	0.28	0.25	0.20	0.20	0.16	0.11	0.09	0.09	0.10	0.11	0.11	0.11
Industrial Processes	0.16	0.16	0.16	0.16	0.16	0.16	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Agriculture	31.83	32.40	32.19	32.74	31.20	31.03	31.18	31.68	31.37	30.72	30.26	29.46	29.11	28.85	29.03
Other ¹	0.40	0.46	0.46	0.46	0.47	0.48	0.47	0.48	0.47	0.53	0.57	0.62	0.69	0.81	0.82
Total	33.05	33.67	33.60	34.20	32.96	32.67	32.61	32.97	32.58	31.91	31.45	30.67	30.36	30.20	30.35

Units: kilotonnes

¹ "Other" for ammonia includes emissions from Energy Industries, Industrial Combustion, Fugitive, Solvent Processes, Waste as well as 1A5b, 7A.


Figure 2-8 Northern Ireland Ammonia Emissions by Sector, 1990-2010



Northern Ireland's ammonia emissions have declined by 8% since 1990 and currently account for 11% of the UK total. The inventory is dominated by emissions from agricultural sources (4) with 79% of the total in 2010 coming from manure management (4B: down 4% since 1990). In 2010, 61% of the Northern Ireland total is from cattle manure management alone (4B1: up 11% since 1990). Other sources of note include transport emissions (1A3: 1% of the Northern Ireland total in 2010), and waste treatment and disposal (6: 2% of the Northern Ireland total in 2010).

Per capita emissions for Northern Ireland are almost four times the UK average in 2010. This is mostly due to the high emissions from agricultural sources, where Northern Ireland contributes 11% to the total UK emission for this sector, compared with only 3% of the UK population.

Figure 2-9 Map of Ammonia Emissions in Northern Ireland, 2010




2.2 CARBON MONOXIDE

Carbon monoxide (CO) arises from incomplete fuel-combustion and is of concern mainly due to its toxicity and its role in tropospheric ozone formation. In terms of human health, CO combines with haemoglobin in blood, decreasing the uptake of oxygen by the lungs, with symptoms varying from nausea to asphyxiation depending upon the level of exposure.

Across the UK, emissions decreased by 77% between 1990 and 2010. This decline has been driven by reductions in emissions from a number of sources, including road transport (1A3b), agricultural field burning (4F) and the domestic sector (1A4b). The decrease is dominated by the reduction in emissions from the road transport sector (1A3b), caused by the increased use of three-way catalysts in cars; this trend is evident for all DAs.

Figure 2-10 UK Carbon Monoxide Emissions by Constituent Country, 1990-2010



The main sources of CO are outlined below:

- Road Transport (1A3b).** Petrol engines are the main source of CO emissions, especially from cold start engine cycles. Since 1990, emissions from road transport sources have reduced by around 83% due to the development of more efficient engine combustion technology, the increased use of catalytic converters and the growth in diesel engine use. Between 2008 and 2009 emissions from passenger cars decreased by 27% due to an improvement in catalyst replacement rates, coupled with a downturn in vehicle km travelled.
- Non road mobile machinery (1A2fii, 1A4bii, 1A4cii).** In the UK, around 14% of CO emissions arise from off-road mobile industrial, residential and agricultural machinery such as portable generators, forklift trucks, lawnmowers and tractors. Emissions are calculated using a complex model which takes into account the population of machinery in the UK, annual usage, the engine size, replacement rates, and the implementation of legislation aimed at reducing emissions. The population of machinery within the model is based on an in depth survey for a single year, with the time series generated using proxy statistics. As such, these estimates are relatively uncertain, however a recent comparison of the UK's NRMM emissions with those of other EU Member States indicated that the emissions are within the expected range for the UK.

- **Stationary domestic combustion (1A4bi).** Around 15% of CO emissions in 2010 arise from residential combustion sources, namely non-electric domestic heating. Reductions in emissions have been significant, due to the switch from solid fuels to the use of gas and electricity, with a 69% reduction in UK emissions since 1990.

Other sources of CO emissions are small compared with transport and off-road sources. Industrial combustion emissions decreased by 26% between 1990 and 2009, reflecting fuel switching from solid fuels to gas, similar to the domestic sector (1A4b). The increase between 2009 and 2010 of 11% reflects the cold weather experienced in 2010. The sudden decline in emissions from the agricultural sector reflects the banning of stubble burning in 1993 in England and Wales, whilst power generation (1A1a) accounts for only 3% of UK emissions in 2010.

Emissions of CO for England, Wales, Scotland and Northern Ireland are summarised in the tables and graphs below. **Table 2-7** shows how total UK CO emissions are split between the four constituent countries, and **Table 2-8** shows per capita emissions for 1990 and 2010.

Table 2-7 Proportion of UK Carbon Monoxide Emissions by Constituent Country

Year	England	Scotland	Wales	Northern Ireland	Unallocated
1990	79%	9%	7%	4%	0%
2010	77%	9%	9%	4%	1%

Table 2-8 Emissions of Carbon Monoxide per capita by Constituent Country (kg/head)

Year	England	Scotland	Wales	Northern Ireland	UK
1990	150.9	159.6	235.5	233.2	158.8
2010	31.5	36.1	63.2	42.9	34.1

2.2.1 England Carbon Monoxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the CO emissions in England by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-9 England Emissions of Carbon Monoxide by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	98.9	92.6	56.3	50.6	59.6	58.0	57.3	63.5	61.7	67.3	67.1	69.0	65.3	57.5	58.4
Industrial Combustion	329.8	349.1	333.4	320.6	298.6	290.7	290.5	289.0	299.2	272.9	272.4	279.5	289.0	255.1	282.7
Transport Sources	5405.5	4595.8	4141.7	3839.5	3351.5	3035.1	2713.3	2425.4	2180.6	1916.1	1692.1	1436.8	1293.8	983.9	834.3
Commercial, domestic and agricultural combustion	679.6	533.3	480.7	495.2	424.5	427.4	353.9	318.5	306.3	263.6	252.4	259.0	277.8	277.9	298.6
Fugitive	22.3	12.3	12.1	9.7	9.7	5.5	4.0	6.2	3.8	3.8	4.1	4.0	3.7	3.7	3.2
Industrial Processes	388.9	413.6	378.7	400.2	354.8	406.5	360.9	268.7	260.3	262.2	263.6	259.3	249.4	192.3	138.5
Other ¹	272.9	32.5	31.0	31.5	30.6	47.0	31.5	32.2	31.0	30.2	30.8	32.0	30.9	29.7	29.6
Total	7198.0	6029.2	5434.0	5147.3	4529.5	4270.2	3811.5	3403.4	3142.9	2816.1	2582.5	2339.6	2210.0	1800.2	1645.3

Units: kilotonnes

¹ "Other" for carbon monoxide includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-11 England Carbon Monoxide Emissions by Sector, 1990-2010



England's CO emissions have declined by 77% since 1990 and account for 77% of the UK total. In 2010, 47% of CO emissions in England stem from road transport combustion sources (1A3bi-iv: down by 86% since 1990), whilst 17% stem from industrial combustion (1A2: down 14% since 1990) and 18% from commercial and residential combustion (1A4: down 56% since 1990). Notable increasing trends in emissions arise from the non-road transport sources, such as railways (1A3c: up by 49% since 1990). However, the levels of emissions from these sources are small relative to emissions that arise from road transport (1A3b) sources.

Between 2008 and 2009, emissions of CO in England have declined by 19%. Much of this change (74%) is driven by the decline in emissions from passenger cars (due to improved catalyst replacement rates), whilst declining emissions from industrial off road machinery and iron and steel have also made a contribution, due to the economic downturn.

Figure 2-12 Map of Carbon Monoxide Emissions in England, 2010



2.2.2 Scotland Carbon Monoxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the CO emissions in Scotland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-10 Scotland Emissions of Carbon Monoxide by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	14.9	15.0	10.4	9.7	11.5	11.2	10.8	9.9	10.4	10.3	12.2	11.7	15.0	14.2	14.1
Industrial Combustion	35.2	32.0	30.1	29.0	27.2	26.9	29.0	28.5	30.1	28.3	27.5	28.2	30.3	25.6	28.2
Transport Sources	507.9	413.2	353.3	323.8	280.9	252.4	226.0	203.5	184.9	163.9	149.1	127.4	115.8	82.2	70.5
Commercial, domestic and agricultural combustion	187.0	121.1	107.9	109.3	94.4	91.8	77.9	69.3	63.1	53.8	55.7	57.1	62.5	62.2	67.1
Fugitive	5.1	1.0	0.9	1.1	1.0	1.7	1.1	0.8	0.8	0.9	0.7	0.9	0.8	0.8	0.8
Industrial Processes	50.3	5.4	5.8	6.7	6.9	7.0	7.2	3.7	3.7	3.6	4.3	4.5	4.6	4.0	4.7
Other ¹	10.3	3.6	3.4	3.4	3.3	7.5	3.4	3.5	3.4	3.3	3.4	3.5	3.4	3.3	3.2
Total	810.8	591.3	511.8	483.0	425.3	398.5	355.4	319.2	296.4	264.1	253.0	233.1	232.4	192.3	188.7

Units: kilotonnes

¹ "Other" for carbon monoxide includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-13 Scotland Carbon Monoxide Emissions by Sector, 1990-2010



Scotland's CO emissions have declined by 77% since 1990 and account for 9% of the UK total. In 2010, 33% of CO emissions in Scotland stem from road transport combustion sources (1A3bi-iv: down by 87% since 1990), whilst 15% stem from industrial combustion (1A2: down 20% since 1990) and 36% from commercial and residential combustion (1A4: down 64% since 1990).

Between 2008 and 2009, emissions fell sharply by 17%, driven predominantly by the reduction in emissions from road transport, with further reductions in emissions from industrial combustion emissions (1A2fi and 1A2fii), driven by the economic downturn.

Figure 2-14 Map of Carbon Monoxide Emissions in Scotland, 2010



2.2.3 Wales Carbon Monoxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the CO emissions in Wales by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-11 Wales Emissions of Carbon Monoxide by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	6.3	6.2	4.4	3.9	5.2	5.9	4.9	5.1	6.8	6.2	6.8	5.4	6.5	5.7	6.6
Industrial Combustion	41.7	30.3	31.5	31.0	26.9	26.8	24.9	25.1	25.0	26.0	27.7	26.9	24.1	19.8	21.8
Transport Sources	322.1	277.2	244.6	226.8	201.8	182.6	164.1	146.1	130.2	113.4	100.9	86.3	77.8	58.6	50.0
Commercial, domestic and agricultural combustion	108.4	89.6	79.2	82.8	68.1	69.2	53.7	46.7	44.5	35.8	36.3	39.4	42.3	43.1	45.3
Fugitive	12.3	19.0	19.0	14.7	15.1	8.5	3.5	4.9	3.3	2.9	6.4	6.9	5.1	5.1	6.2
Industrial Processes	165.7	197.8	206.9	198.7	164.6	143.6	78.6	81.0	103.4	107.6	140.3	124.0	101.1	87.5	58.8
Other ¹	17.4	1.7	1.6	1.6	1.6	2.8	1.6	1.6	1.6	1.5	1.5	1.6	1.5	1.5	1.4
Total	673.9	621.8	587.3	559.5	483.2	439.3	331.3	310.5	314.8	293.4	319.8	290.5	258.4	221.2	190.0

Units: kilotonnes

¹ "Other" for carbon monoxide includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-15 Wales Carbon Monoxide Emissions by Sector, 1990-2010




Wales CO emissions have declined by 72% since 1990 and account for 9% of the UK total. The iron & steel industry (2C1 and 1A2a) contributes significantly to the Welsh total, with a total of 33%. Emissions from iron and steel production have decreased by 65% since 1990, including a 22% reduction between 2009 and 2010. In 2010, 24% of CO emissions in Wales stem from road transport combustion sources (1A3bi-iv: down by 86% since 1990), whilst 24% stem from commercial and residential combustion (1A4: down 58% since 1990). Total CO emissions increased between 2005 and 2006, as a result of additional emissions from the iron & steel industry (1A2), solid fuel transformation (1B1b) and metal production (2C). This increase in emissions was

attributed to additional industrial output from Wales during 2006, but more recent trends show a decline in industrial emissions since 2006.

On a per capita basis, Welsh emissions are higher than the UK average, owing to the large contribution of emissions from iron and steel production. The continuing contribution of emissions from this source has also meant that the percentage decline in emissions per capita in Wales is lower than the other DAs (-73%, compared with a decline of 77-82%).

Figure 2-16 Map of Carbon Monoxide Emissions in Wales, 2010



2.2.4 Northern Ireland Carbon Monoxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the CO emissions in Northern Ireland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-12 Northern Ireland Carbon Monoxide Emissions by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	4.1	3.8	2.1	1.3	1.3	1.4	1.1	1.1	2.0	3.3	2.8	2.6	2.5	2.3	2.0
Industrial Combustion	10.3	11.4	10.7	10.5	9.0	8.4	8.3	8.2	8.5	8.6	9.5	10.2	10.2	8.2	9.2
Transport Sources	175.5	133.9	107.2	99.4	88.5	80.9	71.1	66.6	62.9	57.1	51.3	45.9	43.1	32.3	28.0
Commercial, domestic and agricultural combustion	173.2	121.9	101.9	92.8	83.9	72.3	66.4	55.6	44.4	36.8	34.3	32.5	35.5	34.4	37.1
Fugitive	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Industrial Processes	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other ¹	8.9	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	0.9
Total	372.2	272.2	223.2	205.3	184.0	164.4	148.3	132.5	118.9	106.8	98.9	92.3	92.3	78.2	77.3

Units: kilotonnes

¹ "Other" for carbon monoxide includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-17 Northern Ireland Carbon Monoxide Emissions by Sector, 1990-2010



Northern Ireland's CO emissions have declined by 79% since 1990 and accounted for 4% of the UK total in 2009. 34% of CO emissions in Northern Ireland stem from road transport combustion sources (1A3bi-iv: down by 85% since 1990). In 2010, 48% of the Northern Ireland total emission comes from commercial, agricultural and residential combustion (1A4: down 79% since 1990), which is a much higher contribution than in other DAs (commercial and residential emissions contribute 18%, 36% and 24% within England, Scotland and Wales respectively) due to the greater use of solid fuels as a result of the less well developed gas network in Northern Ireland.

Between 2008 and 2009, emissions fell by 15%, mostly down to the reduction in emissions from passenger cars, but with further significant reductions from industrial (1A2f) and residential combustion (1A4b).

Figure 2-18 Map of Carbon Monoxide Emissions in Northern Ireland, 2010



2.3 NITROGEN OXIDES


Across the UK, emissions of oxides of nitrogen (NO_x) arise primarily from combustion sources. The estimation of these emissions is complex since the nitrogen can be derived from either the fuel or atmospheric nitrogen. The emission is dependent on the conditions of combustion, in particular temperature and excess air ratio, which can vary considerably. Thus combustion conditions, load and even state of maintenance are important. The main three combustion sources of NO_x are:

- **Transport (1A3).** In 2010 road vehicles contributed 34% of total UK NO_x emissions. Since 1990 there has been a steady decline in emissions due to the introduction of catalytic converters on cars and stricter regulations on truck emissions. Between 2008 and 2009, there was about a 21% reduction in the emissions from passenger cars which was mainly driven by improvement in catalyst repair rates⁵. This was due to the introduction of the Regulations Controlling Sale and Installation of Replacement Catalytic Converters and Particle Filters for Light Vehicles for Euro 3 (or above) LDVs after June 2009. However, roadside measurements of ambient NO_x concentrations and a recent study undertaken by King's College London and AEA (Carslaw *et al.*, 2011) have indicated that the trend of ambient NO_x concentration has been fairly flat since 2002, and there is also some evidence from roadside remote sensing of exhaust plumes from a large number of vehicles indicating that NO_x emissions are significantly higher than indicated by the current emissions factors. Consideration is therefore currently underway by Defra and the inventory team to update the NO_x emission factors for the next inventory cycle so that they are more representative of the real world observations and the latest evidence on vehicle emissions. Research indicates that conurbations and city centres show high localised emissions due to the combination of road transport (1A3b), residential and commercial combustion sources. Similarly, around airports, ports and major terminals, significant localised emissions arise from aviation, shipping, railway locomotives and road vehicles.
- **Power Generation (1A1a).** Since 1988 the electricity generators have adopted a programme of progressively fitting low-NO_x burners to their 500 MWe (megawatt electric) or larger coal fired units. More recently the increased use of nuclear generation and the introduction of CCGT (Combined Cycle Gas Turbine) plant burning natural gas have further reduced NO_x emissions. The emissions from the low-NO_x turbines used are much lower than those of pulverised coal fired plant even when low-NO_x burners are fitted. Assuming that these trends continue, power station emissions are expected to fall further. An additional factor has been the recent retrofitting of Boosted Over Fire Air (BOFA) systems to reduce NO_x formation and ensure compliance with the Large Combustion Plant Directive. Between 2008 and 2009, there was an 11% decrease in the emissions from coal burning power stations due to the use of BOFA, as well as the decreased consumption of coal. Emissions decreased by a further 7% between 2009 and 2010.
- **Industrial Combustion (1A2).** The emissions from industrial combustion have declined by 72% since 1970 and they currently contribute 16% to total UK emissions. This is primarily due to the decline in coal use in favour of gas and electricity.

As can be seen in Figure 2.19, total UK emissions of NO_x have decreased since 1990 (by 62%).

⁵ A sensitive parameter in the emission calculations for petrol cars is the assumption made about the proportion of the fleet with catalyst systems that have failed, for example due to mechanical damage or failure of the lambda sensor. Following discussions with DfT, it is assumed that the failure rate is 5% per annum for all Euro standards and that up to 2008, only 20% of failed catalysts were rectified properly, but those that were rectified were done so within a year of failing. The revisions are based on evidence on fitting of replacement catalysts. According to DfT there is evidence that a high proportion of replacement catalysts were not Type Approved and do not restore the emission performance of the vehicle to its original level (DfT 2009). This is being addressed through the Regulations Controlling Sale and Installation of Replacement Catalytic Converters and Particle Filters for Light Vehicles for Euro 3 (or above) LDVs after June 2009. Therefore a change in the repair rate is taken into account for Euro 3 and above petrol LDVs from mid-2009, assuming all failed vehicles are rectified properly.

Figure 2-19 UK Nitrogen Oxides Emissions by Constituent Country, 1990-2010



Emissions of NO_x for England, Wales, Scotland and Northern Ireland are summarised in the tables and graphs below. **Table 2-13** shows how total UK NO_x emissions are split between the four constituent countries and **Table 2-14** shows per capita emissions.

Table 2-13 Proportion of UK Nitrogen Oxides Emissions by Constituent Country

Year	England	Scotland	Wales	Northern Ireland	Unallocated
1990	79%	10%	6%	3%	2%
2010	75%	10%	7%	3%	5%

Table 2-14 Emissions of NO_x per capita by Constituent Country (kg/head)

Year	England	Scotland	Wales	Northern Ireland	UK
1990	47.9	55.2	61.5	61.0	50.4
2010	15.8	21.1	27.2	18.5	17.8

2.3.1 England Nitrogen Oxides Inventory by Sector, 1990-2010

The table and graph below give a summary of the NO_x emissions in England by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-15 England Emissions of Nitrogen Oxides by sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	651.1	416.5	314.5	279.5	294.0	314.4	310.3	344.1	326.9	347.1	338.3	327.3	243.3	210.8	208.2
Industrial Combustion	321.4	298.6	268.0	254.8	243.6	233.5	223.6	216.2	212.3	213.8	194.2	194.0	177.3	143.9	145.2
Transport Sources	1090.7	908.6	782.1	734.4	678.8	651.1	613.5	578.4	558.5	537.6	518.8	496.0	466.8	395.8	373.1
Commercial, domestic and agricultural combustion	154.0	142.2	134.7	130.5	122.1	119.3	107.3	101.8	96.1	90.9	82.5	76.6	73.6	66.5	69.5
Other ¹	68.2	56.6	39.1	41.0	37.5	37.3	32.9	35.1	36.1	33.3	33.1	36.9	35.2	31.8	30.3
Total	2285.3	1822.5	1538.4	1440.2	1376.0	1355.6	1287.8	1275.6	1229.9	1222.8	1167.0	1130.8	996.3	848.7	826.4

Units: kilotonnes


¹ "Other" for nitrogen oxides includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-20 England Nitrogen Oxides Emissions by Sector, 1990-2010



England's NO_x emissions have declined by 64% since 1990 and account for 75% of the UK total. Power generation (1A1a) is a very significant source, accounting for 23% of the England total in 2010, although emissions from this source have reduced by 70% since 1990. In 2010, 37% of NO_x emissions in England stem from road transport combustion sources (1A3bi-iv: down by 70% since 1990), whilst 18% stem from industrial combustion (1A2: down 55% since 1990). Notable increases in emissions arise from railways (1A3c: up by 72% since 1990 accounting for 3% of the 2010 England's total emission) and from international aviation landing and take-off (LTO) (1A3ai(i): up by 96% since 1990 and 1A3a(ii): up by 20% since 1990 in 2010). Aviation emissions account for only 1% of the emissions in England in 2010.

Figure 2-21 Map of Nitrogen Oxides Emissions in England, 2010



2.3.2 Scotland Nitrogen Oxides Inventory by Sector, 1990-2010

The table and graph below give a summary of the NO_x emissions in Scotland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-16 Scotland Emissions of Nitrogen Oxides by sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	96.8	65.5	52.0	49.9	56.3	51.7	50.2	47.2	45.9	45.9	58.4	51.5	40.9	35.3	39.3
Industrial Combustion	32.0	27.7	25.3	24.1	23.8	23.8	20.7	19.6	19.7	20.5	18.9	18.9	18.5	14.8	14.5
Transport Sources	117.4	96.0	81.3	76.9	70.7	67.2	64.3	60.9	58.9	57.4	55.9	54.2	51.2	43.4	41.1
Commercial, domestic and agricultural combustion	26.5	25.1	24.2	23.6	22.2	21.9	20.0	19.0	17.6	16.9	15.0	13.8	13.1	12.1	12.2
Other ¹	7.8	4.0	3.3	3.9	3.8	4.0	3.7	3.7	3.7	3.6	3.4	3.4	3.3	2.9	3.0
Total	280.4	218.3	186.1	178.3	176.8	168.7	159.0	150.4	145.8	144.3	151.6	141.7	127.2	108.4	110.0

Units: kilotonnes

¹ "Other" for nitrogen oxides includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-22 Scotland Nitrogen Oxides Emissions by Sector, 1990-2010




Scotland's NO_x emissions have declined by 61% since 1990 (Figure 2-22) and currently account for 10% of the UK total. Power generation (1A1a) is a very significant source of NO_x emissions, accounting for 29% of the Scotland total in 2010; although emissions from this source have reduced by 63% since 1990. (Note that in the table and figure above, the sector "Energy Industries" includes power generation, petroleum refining and other energy industries such as collieries and gas processing.)

Recent trends in electricity generation have dominated the overall trends in the inventory. Although overall electricity generation in Scotland decreased between 2009 and 2010 (by 3%), coal-fired generation increased by 23% to 14,653 GWh. Gas-fired generation decreased by 14% to 6,618 GWh.

A further 29% of NO_x emissions in Scotland arise from road transport sources (1A3b: down by 70% since 1990), 13% stem from industrial combustion (1A2: down 55% since 1990) and 5% is from agricultural mobile machinery (1A4cii, down 63% since 1990). Increases in emissions are apparent mostly in relatively minor source sectors such as international aviation landing and take-off (LTO) (1A3ai(i): up by 122% since 1990). Emissions from rail have also increased by 109% since 1990, now contributing 3% to the total emissions in Scotland. This is due to increases in fuel oil consumption by the rail sector from 1990 due to rise in passenger train km and freight train km during this time.

Figure 2-23 Map of Nitrogen Oxides Emissions in Scotland, 2010



2.3.3 Wales Nitrogen Oxides Inventory by Sector, 1990-2010

The table and graph below give a summary of the NO_x emissions in Wales by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-17 Wales Emissions of Nitrogen Oxides by sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	50.0	35.4	28.8	23.5	32.6	38.9	33.0	32.7	35.7	34.1	41.4	29.1	39.2	30.4	30.6
Industrial Combustion	30.9	30.5	29.4	31.4	28.2	25.5	20.6	20.9	20.0	19.0	17.7	17.3	15.9	12.7	13.2
Transport Sources	69.6	59.1	51.1	47.8	44.5	42.4	40.4	38.3	37.5	36.1	34.9	33.8	32.4	28.1	26.9
Commercial, domestic and agricultural combustion	17.2	16.6	15.8	15.4	14.5	14.2	13.1	12.4	11.6	10.9	9.7	8.7	8.2	7.5	7.4
Other ¹	8.2	8.7	8.3	8.5	8.2	6.3	2.9	3.3	3.7	3.6	4.0	4.0	4.0	3.0	3.7
Total	175.9	150.2	133.3	126.6	128.0	127.3	110.0	107.5	108.5	103.7	107.8	93.0	99.6	81.6	81.8

Units: kilotonnes

¹ "Other" for nitrogen oxides includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-24 Wales Nitrogen Oxides Emissions by Sector, 1990-2010



Wales NO_x emissions have declined by 54% since 1990 and accounted for 7% of the UK total in 2010. Power generation (1A1a) accounts for 29% of the Wales NO_x inventory total in 2010; emissions from this source have reduced by 44% since 1990. Recent trends in electricity generation have dominated the overall trends in the Wales NO_x inventory, with large fluctuations in coal-fired generation in particular (due to a plant shut-down at Aberthaw during 2007 to retro-fit 2 units with Flue Gas Desulphurisation abatement). Between 2007 and 2008, coal-fired generation increased by nearly 83% (up to 9,364 GWh) once Aberthaw came back on-line, then decreasing between 2008 and 2009 by 30% (down to 6,547 GWh). Gas-fired generation has increased by 6% between 2009 and 2010 to 15,220 GWh, whilst coal-fired generation has decreased by 9% to 5,929 GWh. As a result, the overall NO_x emissions from the sector have remained relatively stable, increasing by 1% between 2009 and 2010 (Figure 2-24).

A further 24% of NO_x emissions in Wales stem from road transport combustion sources (1A3bi-iv: down by 70% since 1990), 16% stem from industrial combustion (1A2: down 57% since 1990) and 5% of emissions are from petroleum refining (1A1b: down 28% since 1990). Notable increases in emissions arise from railways (1A3c: up by 237% since 1990 to 5% of the 2010 Wales total) and from aviation landing and take-off (LTO) (1A3ai(i): up by 96% since 1990 and 1A3aii(i): up by 78% since 1990 in 2010; combined, these sources account for less than 1% of the emissions in Wales in 2010).

Figure 2-25 Map of Nitrogen Oxides Emissions in Wales, 2010



2.3.4 Northern Ireland Nitrogen Oxides Inventory by Sector, 1990-2010

The table and graph below give a summary of the NO_x emissions in Northern Ireland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-18 Northern Ireland Emissions of Nitrogen Oxides by sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	31.0	19.3	13.9	14.2	14.8	16.0	12.3	11.5	9.8	9.6	9.6	7.0	6.4	5.0	5.7
Industrial Combustion	13.4	12.3	10.3	9.3	8.5	8.4	7.5	7.1	7.1	8.4	7.7	8.2	7.7	5.3	5.3
Transport Sources	37.0	28.0	21.9	21.1	20.1	19.6	19.9	20.6	19.7	19.5	18.9	18.5	17.7	14.9	14.1
Commercial, domestic and agricultural combustion	13.9	13.0	12.9	12.7	12.2	12.0	11.6	11.2	10.4	10.1	9.2	8.2	7.8	7.5	7.5
Other ¹	2.0	1.2	1.0	1.1	1.0	0.9	0.8	0.8	0.8	0.8	0.8	0.9	0.8	0.8	0.7
Total	97.4	73.8	60.0	58.4	56.7	57.0	52.2	51.2	47.8	48.4	46.3	42.7	40.5	33.5	33.3

Units: kilotonnes

¹ "Other" for nitrogen oxides includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-26 Northern Ireland Nitrogen Oxides Emissions by Sector, 1990-2010



Northern Ireland's NO_x emissions have declined by 66% since 1990 and account for only 3% of the UK total. 36% of NO_x emissions in Northern Ireland stem from road transport combustion sources (1A3bi-iv: down by 66% since 1990), whilst 16% stem from industrial combustion (1A2: down 60% since 1990). Power generation (1A1a) accounted for 17% of the Northern Irish total in 2010 and emissions from this source have reduced by 82% since 1990. 10% are from agricultural off road machinery (1A4cii: down 62% since 1990).

Notable increases in emissions arise from very minor sources sector such as international aviation LTO (1A3ai(i): up by 235% since 1990). However, this source accounts for less than 1% of the Northern Ireland total emission in 2010.

Figure 2-27 Map of Nitrogen Oxides Emissions in Northern Ireland, 2010



2.4 NON-METHANE VOLATILE ORGANIC COMPOUNDS

Non-Methane Volatile Organic Compounds (NMVOCs) are emitted to air as combustion products, as vapour arising from handling or use of petroleum distillates, solvents or chemicals, and from numerous other sources. The diversity of processes which emit NMVOCs is huge, covering not only many branches of industry, but also transport (1A3), agriculture (4) and domestic sources.

UK emissions inventory data indicate that only 19% of the NMVOC emissions arise from combustion sources (unlike SO₂ and NO_x where the contribution from combustion sources is much higher with 92% and 99% respectively). Of these emissions from combustion sources, it is the transport (1A3) sector that dominates (70% of combustion emissions). NMVOC emissions are dependent on vehicle speed and are higher on minor and urban major roads than on the high-speed motorways and major roads.

A large proportion of emissions are caused either as a result of the activities of people in and around their homes (e.g. domestic solvent use or domestic combustion), or by widespread industrial activities such as small-scale industrial coating processes, dry cleaners and small bakeries.


- **Solvent and other product use (3).** This sector comprises industrial and domestic solvent applications (cleaning, degreasing), as well as the manufacturing and processing of chemical products. It represents 44% of the UK total NMVOC emission in 2010. During the 1990s, industrial NMVOC emissions have fallen as a result of emission controls, technological changes, and reduced manufacturing output in some sectors. Emissions from the chemical industry have reduced during the 1990s as tighter emission controls have been introduced. Domestic solvent emissions have also fallen due to a trend towards formulating products such as paints and aerosols with lower solvent contents.
- **Stationary Combustion.** This sector includes emissions from public electricity and heat production as well as those from petroleum refining and the manufacture of iron and steel. Emissions from the petroleum-refineries have fallen significantly due to a reduction in refinery capacity and tighter emission regulations during the 1990s.
- **Production processes.** This sector includes emissions from metal production, road construction, and non-fuel mining. These processes are estimated to comprise approximately 3% of the UK total emission in 2010.
- **Processes in wood, paper pulp and food & drink.** Emissions from the food and drink industry comprised approximately 11% of the total NMVOC emission in 2010. The largest source is whisky maturation although bread baking, animal feed manufacture, fat and oil processing and barley malting are also important sources.
- **Transport (1A3).** Emissions from transport sources are currently responsible for 9% of NMVOC emissions, with road transport (1A3b) accounting for most of this. During the 1990s, these emissions have declined significantly due to the increased use of catalytic converters and fuel switching from petrol to diesel cars. Between 2008 and 2009, emissions from passenger cars fell by 32% due to the assumed improvements in catalyst repair rates.
- **Offshore oil and gas.** Emissions from this sector have increased substantially with the growth of the UK's offshore activities, and stem primarily from tanker loading / unloading.

Other sources of NMVOCs include:

- Gas leakage from the national gas distribution networks;
- Evaporative losses from the distribution and marketing of petrol;
- Waste treatment and disposal contribute;
- Natural and agricultural sources.

UK emissions of NMVOC fell by 71% between 1990 and 2010, primarily due to reductions in road transport sources through the use of catalytic converters and fuel switching to diesel technology.

Figure 2-28 UK NMVOC Emissions by Constituent Country, 1990-2010



Emissions of NMVOCs for England, Wales, Scotland and Northern Ireland are summarised in the tables and graphs below. **Table 2-19** shows how total UK NMVOC emissions are split between the 4 constituent countries and **Table 2-20** shows per capita emissions.

Table 2-19 Proportion of UK NMVOC Emissions by Constituent Country

Year	England	Scotland	Wales	Northern Ireland	Unallocated
1990	74%	15%	5%	2%	3%
2010	72%	16%	5%	2%	5%

Table 2-20 Emissions of NMVOC per capita by Constituent Country (kg/head)

Year	England	Scotland	Wales	Northern Ireland	UK
1990	43.1	80.2	48.9	41.7	48.2
2010	10.9	23.4	14.4	10.5	12.7

2.4.1 England NMVOC Inventory by Sector, 1990-2010

The table and graph below give a summary of the NMVOC emissions in England by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-21 England Emissions of NMVOC by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	5.8	6.2	3.6	3.3	4.7	4.0	5.6	4.1	4.1	3.8	4.5	3.0	3.1	2.9	2.9
Industrial Combustion	24.5	25.1	25.3	24.9	24.6	24.8	24.2	23.2	23.4	24.2	22.3	23.3	22.0	18.1	18.8
Transport Sources	856.6	673.3	539.4	481.8	411.1	352.2	298.6	250.9	212.1	179.3	154.6	130.0	115.4	82.2	69.9
Commercial, domestic and agricultural combustion	57.7	42.1	42.8	44.5	37.8	35.6	32.2	31.2	29.9	27.9	26.5	25.9	26.4	24.7	25.2
Fugitive	270.7	230.2	216.1	193.2	182.7	187.0	166.4	140.8	133.9	129.8	117.7	112.9	101.2	95.1	86.1
Industrial Processes	184.4	167.9	120.9	89.1	85.9	74.9	73.9	79.0	62.5	59.7	53.5	52.2	46.0	40.4	40.6
Solvent Processes	557.3	450.5	416.3	393.4	372.2	359.8	353.4	348.6	349.0	343.4	342.8	337.0	318.7	296.4	294.9
Other ¹	97.5	65.5	56.1	52.1	49.2	43.9	40.8	36.7	34.1	33.2	32.9	32.5	31.5	30.4	29.4
Total	2054.5	1660.8	1420.6	1282.4	1168.3	1082.1	995.1	914.6	848.9	801.1	754.9	716.9	664.2	590.1	568.0

Units: kilotonnes

¹ "Other" for NMVOC includes emissions from Agriculture, Waste as well as 1A5b, 7A.


Figure 2-29 England NMVOC Emissions by Sector, 1990-2010



England's NMVOC emissions have declined by 72% since 1990 and account for 72% of the UK total. The solvent and other product use sector (3) is a very significant source, accounting for 52% of the total in 2010. Emissions from this sector have reduced by 47% since 1990. Fugitive emissions from fuels (1B) account for

15% of the total in 2010 with emissions down 68% since 1990. Further significant sources are road transport, including evaporative losses (1A3bi-v), these are 11% of the total in 2010; emissions from this source have reduced by 93% since 1990 and industrial processes (2), which accounts for 7% of the total in 2010, with emissions down 78% since 1990.

Figure 2-30 Map of NMVOC Emissions in England, 2010



2.4.2 Scotland NMVOC Inventory by Sector, 1990-2010

The table and graph below give a summary of the NMVOC emissions in Scotland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-22 Scotland Emissions of NMVOC by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	0.9	1.0	0.7	0.4	0.7	0.6	0.7	0.6	0.6	0.6	0.8	0.5	0.5	0.5	0.5
Industrial Combustion	2.4	2.4	2.4	2.4	2.4	2.4	2.3	2.2	2.2	2.2	2.1	2.2	2.1	1.7	1.8
Transport Sources	79.5	59.5	44.8	39.4	33.3	28.3	24.1	20.3	17.4	15.0	13.3	11.4	10.2	6.5	5.6
Commercial, domestic and agricultural combustion	13.0	8.3	8.3	8.6	7.2	6.7	6.0	5.8	5.4	5.1	5.3	5.3	5.5	5.3	5.5
Fugitive	176.9	117.0	97.8	81.7	79.0	68.4	71.6	52.5	39.2	40.3	38.2	38.5	32.1	18.8	17.3
Industrial Processes	61.6	61.4	60.9	62.5	61.9	60.8	60.0	60.9	58.0	56.9	56.9	58.5	57.9	58.3	58.2
Solvent Processes	58.6	47.8	44.4	38.2	36.0	34.8	33.7	33.3	33.3	32.5	32.4	32.1	30.6	28.7	28.7
Other ¹	14.4	11.9	9.9	9.1	8.5	7.4	6.8	6.1	5.6	5.5	5.4	5.3	5.1	5.0	4.8
Total	407.4	309.2	269.3	242.3	229.1	209.4	205.3	181.7	161.8	158.0	154.3	153.7	144.1	124.8	122.4

Units: kilotonnes

¹ "Other" for NMVOC includes emissions from Agriculture, Waste as well as 1A5b, 7A.

Figure 2-31 Scotland NMVOC Emissions by Sector, 1990-2010




Scotland's NMVOC emissions have declined by 70% since 1990 and account for 16% of the UK total. Industrial processes (2) are a significant source making up 48% of the total in 2010, with emissions down 6% since 1990. Industrial processes include food and drink emissions (2D2) dominated by brewers and distilleries.

This is 44% of the Scottish total in 2010, up 23% since 1990. Other significant sources are solvents and other product use (3) which account for 23% of the total in 2010. Emissions from this source are down 51% since 1990. Fugitive emissions from fuels (1B) make up 14% of the total in 2010, these emissions have reduced by 90% since 1990. Road transport is also a significant source, including evaporative losses (1A3b) accounting for 4% of the total in 2010, with emissions down 94% since 1990.

On a per capita basis, Scottish emissions are around 85% higher than the UK average, due to the high contribution of NMVOC emissions from food and drink processes.

Figure 2-32 Map of NMVOC Emissions in Scotland, 2010



2.4.3 Wales NMVOC Inventory by Sector, 1990-2010

The table and graph below give a summary of the NMVOC emissions in Wales by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-23 Wales Emissions of NMVOC by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	1.0	1.1	0.7	0.5	0.7	0.5	0.7	0.6	0.5	0.4	0.4	0.3	0.4	0.3	0.4
Industrial Combustion	1.9	2.0	2.0	2.1	2.2	2.1	2.0	2.0	1.9	1.8	1.7	1.8	1.6	1.4	1.4
Transport Sources	50.4	40.6	31.9	28.4	24.7	21.2	18.1	15.1	12.7	10.6	9.2	7.8	6.9	4.9	4.2
Commercial, domestic and agricultural combustion	8.1	5.9	6.1	6.5	5.1	4.7	4.0	3.8	3.6	3.4	3.5	3.7	3.9	3.7	3.8
Fugitive	22.8	21.1	17.4	14.5	15.5	12.7	13.0	11.8	12.5	11.6	12.3	13.0	11.5	12.0	11.7
Industrial Processes	5.5	5.2	5.0	4.6	4.1	3.3	2.2	5.4	4.2	2.3	2.5	2.4	2.2	2.1	2.2
Solvent Processes	43.7	34.2	28.8	27.5	23.2	21.8	21.2	20.8	20.8	21.2	21.3	20.8	19.3	17.8	17.6
Other ¹	6.3	4.2	3.6	3.4	3.2	2.8	2.6	2.3	2.1	2.1	2.1	2.0	2.0	1.9	1.9
Total	139.8	114.3	95.5	87.5	78.8	69.2	63.8	61.8	58.4	53.3	52.9	51.8	47.8	44.1	43.2

Units: kilotonnes

¹ "Other" for NMVOC includes emissions from Agriculture, Waste as well as 1A5b, 7A.


Figure 2-33 Wales NMVOC Emissions by Sector, 1990-2010



Wales' NMVOC emissions have declined by 69% since 1990 and account for 5% of the UK total. Solvents and other product use (3) accounts for 41% of the Wales NMVOC inventory total in 2010; emissions from this source have reduced by 60% since 1990. Fugitive emissions from fuels (1B) are another significant source of emissions making up 27% of the total in 2010. Emissions from this source have been reduced by 49%

since 1990. Road transport is an important source of emissions, including evaporative losses (1A3bi-v), accounting for 8% of the total in 2010. Emissions from this source are down 93% since 1990. Industrial processes (2) account for 5% of the total in 2010, with emissions down 60% since 1990.

Figure 2-34 Map of NMVOC Emissions in Wales, 2010



2.4.4 Northern Ireland NMVOC Inventory by Sector, 1990-2010

The table and graph below give a summary of the NMVOC emissions in Northern Ireland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-24 Northern Ireland Emissions of NMVOC by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	0.3	0.4	0.2	0.2	0.2	0.0	0.0	0.0	0.2	0.3	0.4	0.1	0.2	0.2	0.2
Industrial Combustion	0.9	0.9	0.9	0.9	0.8	0.8	0.8	0.8	0.8	0.9	0.8	0.9	0.8	0.7	0.7
Transport Sources	28.6	19.2	13.1	11.6	10.0	8.7	7.2	6.4	5.7	4.9	4.3	3.8	3.5	2.2	1.9
Commercial, domestic and agricultural combustion	10.5	7.0	6.1	5.7	5.2	4.7	4.4	3.9	3.3	3.0	2.8	2.6	2.6	2.5	2.6
Fugitive	3.4	3.2	2.9	2.2	2.1	1.9	1.8	1.7	1.7	1.5	1.4	1.3	1.1	1.0	0.9
Industrial Processes	2.4	2.4	2.2	2.1	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.1	2.1	2.1	2.1
Solvent Processes	17.4	13.8	12.8	12.2	11.9	11.6	11.4	11.2	11.2	11.0	11.1	10.9	10.4	9.7	9.6
Other ¹	3.1	2.1	1.8	1.7	1.6	1.4	1.3	1.2	1.1	1.1	1.1	1.1	1.0	1.0	1.0
Total	66.6	48.9	39.9	36.5	33.9	31.2	29.0	27.3	25.9	24.8	23.9	22.7	21.8	19.4	19.0

Units: kilotonnes

¹ "Other" for NMVOC includes emissions from Agriculture, Waste as well as 1A5b, 7A.


Figure 2-35 Northern Ireland NMVOC Emissions by Sector, 1990-2010



Northern Ireland's NMVOC emissions have declined by 71% since 1990 and account for 2% of the UK total. Solvent and other product use (3) are a major source making up 51% of the NMVOC inventory total in Northern Ireland in 2010. Emissions from this source are down 45% since 1990. Another significant source is commercial & domestic combustion (1A4), which accounts for 14% of the NMVOC inventory total in Northern

Ireland in 2010. Emissions from this source are down 76% since 1990. The food & drink sector (2D2) accounts for 11% of the total in 2010 and emissions from this source have increased by 12% since 1990 due to increased emissions from whisky production. Road transport sources, including evaporative losses (1A3bi-v) make up 9% of the total in 2010 with a 94% reduction since 1990. Another important source is fugitive emissions from fuels (1B) accounting for 5% of the total in 2010. This source of emission has reduced by 75% since 1990.

Figure 2-36 Map of NMVOC Emissions in Northern Ireland, 2010



2.5 PARTICULATE MATTER AS PM₁₀

PM₁₀ is a measure of the size distribution of the particles emitted to air and represents the proportion material with an aerodynamic diameter less than 10 micro meters. PM₁₀ in the atmosphere arises from primary and secondary sources:

Primary Sources

Direct emissions of particulate matter into the atmosphere arise from a wide range of sources such as fuel combustion, surface erosion and wind-blown dusts and mechanical break-up in, for example, quarrying and construction sites.

Secondary Sources

Particulate matter may be formed in the atmosphere through reactions of other pollutants such as sulphur dioxide, nitrogen oxides and ammonia to form solid sulphates and nitrates, as well as organic aerosols formed from the oxidation of NMVOCs.

These inventories only consider primary sources. For further information on secondary particulates see the Air Quality Expert Group's Report on particulate matter in the United Kingdom (AQEG, 2005).


The main sources of primary PM₁₀ are briefly described below:

- **Road Transport (1A3b).** Diesel engines typically emit a greater mass of particulates per vehicle kilometre than petrol engines. Particulate emissions also arise from all vehicles through brake and tyre wear as well as from the re-entrainment of dust from road surfaces caused by vehicle movements.
- **Stationary Combustion (1A1, 1A2 and 1A4).** Domestic coal combustion has historically been the main source of particulate emissions in the UK, but restrictions in the use of coal for domestic combustion through the Clean Air Acts has led to other sources becoming more important nationally. Domestic coal is still a significant source in Northern Ireland, some smaller towns and villages, and in areas associated with the coal industry. Other fossil fuels emit PM₁₀ with combustion of wood, gas oil and fuel oil all contributing significantly to UK emissions. In general, particles emitted from fuel combustion are of a smaller size than from other sources.
- **Industrial Processes (2).** Particulates are emitted from a wide range of industrial processes including: the production of metals, cement, lime, coke & chemicals, bulk handling of dusty materials, construction, mining and quarrying. Whilst emission monitoring results are now widely available for stack and other point-source emissions of particulates from regulated industrial processes, the quantification of diffuse & fugitive emissions from industrial sources is more difficult. Few UK measurements are available for these fugitive releases but there have been substantial improvements in the estimation of PM₁₀ emissions from industrial processes in recent years.

2.5.1 UK Trends in PM₁₀ Emissions

Emissions of PM₁₀ from across the UK have declined significantly since 1970, mainly due to improved abatement of industrial and power generation (1A1a) emission sources and a general reduction in coal use as an energy source across many economic sectors. For example, emissions in the domestic and commercial sector have fallen from 246 kt (49% of the total emission) in 1970 to 19 kt (16%) in 2010.

Figure 2-37 UK PM₁₀ Emissions by Constituent Country, 1990-2010



Emissions from power stations have declined despite a significant growth in electricity generation capacity, due to a shift in the fuel mix for power generation from coal to natural gas, nuclear and renewable generation, and also due to abatement being fitted at coal-fired power stations. For example, the installation of flue gas desulphurisation (FGD) at a number of plants has reduced particulate matter emissions substantially. Emissions from road transport have varied across the time-series as a number of factors have combined. The main source of road transport emissions is exhaust gases from diesel engines. Emissions from diesel vehicles have been growing due to the growth in heavy-duty vehicle traffic and the move towards more diesel cars. Since around 1992, however, emissions from diesel vehicles have been decreasing due to the penetration of new vehicles meeting tighter PM₁₀ emission regulations ("Euro standards" for diesel vehicles were first introduced in 1992).

Among the non-combustion and non-transport sources, the major emissions are from industrial processes (2), the most important of which is quarrying whose emission rates have remained fairly constant. Other industrial processes, including the manufacture of steel, cement, lime, coke, and primary and secondary non-ferrous metals, are collectively important sources of particulate matter although emissions from individual sectors are relatively insignificant.

Emissions of PM₁₀ for England, Wales, Scotland and Northern Ireland are summarised in the tables and graphs below. **Table 2-25** shows how total UK PM₁₀ emissions are split between the 4 constituent countries, and **Table 2-26** show per capita emissions.

Table 2-25 Proportion of UK PM₁₀ Emissions by Constituent Country

Year	England	Scotland	Wales	Northern Ireland	Unallocated
1990	76%	11%	7%	6%	1%
2010	75%	11%	8%	5%	1%

Table 2-26 Emissions of PM₁₀ per capita by Constituent Country (kg/head)

Year	England	Scotland	Wales	Northern Ireland	UK
1990	4.5	5.9	7.0	9.9	4.9
2010	1.6	2.5	3.1	2.9	1.8

2.5.2 England PM₁₀ Inventory by Sector, 1990-2010

The table and graph below give a summary of the PM₁₀ emissions in England by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-27 England Emissions of PM₁₀ by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	59.9	34.0	22.4	18.2	19.0	13.4	7.4	7.6	7.8	8.8	9.1	7.4	7.6	6.0	5.6
Industrial Combustion	22.4	21.4	19.8	18.3	16.5	16.6	15.4	14.3	14.2	13.8	11.9	11.6	10.7	9.0	8.9
Transport Sources	37.4	39.7	37.2	36.3	32.3	31.6	30.6	29.8	29.2	28.6	27.8	26.4	25.1	23.8	23.0
Commercial, domestic and agricultural combustion	37.8	23.8	24.5	25.9	20.5	19.0	16.0	15.1	14.6	14.1	14.0	14.5	16.3	16.1	17.2
Fugitive	0.4	0.2	0.3	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.6	0.6	0.5	0.3	0.5
Industrial Processes	30.9	26.7	21.8	20.6	18.9	19.1	18.5	18.8	18.2	18.0	17.2	18.4	16.8	14.0	13.0
Other ¹	23.6	20.6	21.3	21.3	21.1	24.0	20.5	21.1	20.6	20.1	20.5	20.0	19.2	17.4	17.7
Total	212.3	166.4	147.3	140.8	128.6	123.9	108.8	107.0	105.1	103.9	101.0	98.9	96.1	86.6	86.1

Units: kilotonnes


¹ "Other" for PM₁₀ includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-38 England PM₁₀ Emissions by Sector, 1990-2010



England's PM₁₀ emissions have declined by 59% since 1990 and account for 75% of the UK total. 24% of PM₁₀ emissions in England come from road transport (1A3b) sources (down by 39% since 1990), whilst 13% stem from commercial and residential combustion (mainly of coal and solid fuels, down by 63% since 1990). Emissions from power generation (1A1a) were 27% of the England total emission in 1990, but have been significantly reduced to 6% of the England total in 2010.

Figure 2-39 Map of PM₁₀ Emissions in England, 2010



2.5.3 Scotland PM₁₀ Inventory by Sector, 1990-2010

The table and graph below give a summary of the PM₁₀ emissions in Scotland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-28 Scotland Emissions of PM₁₀ by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	8.3	5.2	3.8	3.1	3.6	3.5	2.2	1.2	1.9	1.8	2.7	2.4	1.0	1.1	1.4
Industrial Combustion	2.2	2.0	1.7	1.6	1.6	1.6	1.4	1.3	1.3	1.3	1.2	1.2	1.1	0.9	0.9
Transport Sources	4.4	4.6	4.3	4.2	3.7	3.5	3.4	3.3	3.2	3.2	3.1	2.9	2.7	2.6	2.5
Commercial, domestic and agricultural combustion	9.4	5.6	5.6	5.9	4.8	4.4	3.8	3.6	3.3	3.2	3.6	3.7	4.0	4.0	4.2
Fugitive	0.1	0.0	0.0	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.0	0.5	0.1	0.6
Industrial Processes	3.0	2.3	1.9	1.8	1.7	1.7	1.7	1.7	1.7	1.7	1.6	1.8	1.4	1.1	1.2
Other ¹	2.5	2.5	2.5	2.4	2.5	3.3	2.5	2.4	2.4	2.3	2.3	2.3	2.3	2.1	2.2
Total	29.9	22.2	19.8	19.0	18.0	18.2	15.2	13.6	14.0	13.6	14.5	14.2	13.0	11.9	12.9

Units: kilotonnes


¹ "Other" for PM₁₀ includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-40 Scotland PM₁₀ Emissions by Sector, 1990-2010



Scotland's PM₁₀ emissions have declined by 57% since 1990 and account for 11% of the UK total. 17% of PM₁₀ emissions in Scotland come from road transport (1A3b) sources (down by 40% since 1990), whilst 24% stem from commercial and residential combustion (mainly of coal and solid fuels, down by 59% since 1990). Emissions from power generation (1A1a) were 25% of the Scotland total emission in 1990, but have been reduced to 10% of the Scotland total in 2010.

Figure 2-41 Map of PM₁₀ Emissions in Scotland, 2010



2.5.4 Wales PM₁₀ Inventory by Sector, 1990-2010

The table and graph below give a summary of the PM₁₀ emissions in Wales by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-29 Wales Emissions of PM₁₀ by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	3.3	2.2	1.7	1.3	2.0	1.7	1.0	1.2	1.0	1.1	1.3	1.2	1.1	0.9	0.8
Industrial Combustion	1.9	1.7	1.3	1.4	1.4	1.4	1.2	1.1	1.0	0.9	0.8	0.8	0.7	0.6	0.6
Transport Sources	2.5	2.7	2.7	2.6	2.3	2.2	2.2	2.1	2.1	2.0	2.0	1.8	1.8	1.7	1.6
Commercial, domestic and agricultural combustion	6.1	4.3	4.5	4.8	3.6	3.3	2.7	2.6	2.5	2.3	2.5	2.8	3.0	2.9	3.0
Fugitive	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Industrial Processes	4.5	4.4	3.9	3.7	3.5	2.5	1.9	2.6	2.8	2.5	2.6	2.8	2.2	1.6	1.9
Other ¹	1.5	1.3	1.4	1.5	1.5	1.6	1.2	1.4	1.3	1.2	1.2	1.2	1.2	1.1	1.1
Total	20.0	16.6	15.5	15.3	14.4	12.8	10.3	11.0	10.7	10.1	10.5	10.7	10.1	9.0	9.2

Units: kilotonnes

¹ "Other" for PM₁₀ includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-42 Wales PM₁₀ Emissions by Sector, 1990-2010



Wales PM₁₀ emissions have declined by 54% since 1990 and accounted for 8% of the UK total in 2010. Commercial and domestic sources accounted for 28% of Welsh emissions in 2010, mainly from combustion of solid fuels; these emissions have declined by 48% since 1990. In 2010, 15% of PM₁₀ emissions in Wales come from road transport (1A3b) sources (down by 33% since 1990), whilst emissions from power generation (1A1a) accounted for 4% of the Wales total in 2010, down 85% on 1990 emissions. Heavy industry plays a

more significant role in the Wales PM₁₀ inventory than in other parts of the UK, with key contributions to the 2010 total from quarrying and mining (4%), iron & steel production (14%), other manufacturing combustion (7%). Welsh per capita emissions of PM₁₀ are 68% higher than the UK average, due primarily to emissions from iron and steel manufacture.

Figure 2-43 Map of PM₁₀ Emissions in Wales, 2010



2.5.5 Northern Ireland PM₁₀ Inventory by Sector, 1990-2010

The table and graph below give a summary of the PM₁₀ emissions in Northern Ireland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-30 Northern Ireland Emissions of PM₁₀ by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	2.7	1.5	1.0	0.6	0.8	1.0	0.4	0.4	0.3	0.3	0.3	0.3	0.2	0.1	0.1
Industrial Combustion	1.0	1.0	0.8	0.8	0.8	0.7	0.6	0.5	0.6	0.7	0.6	0.7	0.6	0.5	0.5
Transport Sources	1.2	1.5	1.4	1.4	1.3	1.2	1.3	1.3	1.2	1.2	1.2	1.1	1.1	1.1	1.0
Commercial, domestic and agricultural combustion	8.7	5.7	4.9	4.5	4.2	3.7	3.4	3.0	2.5	2.2	2.1	1.9	2.0	1.9	2.0
Fugitive	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Industrial Processes	0.7	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.3	0.3
Other ¹	1.5	1.6	1.6	1.6	1.5	1.4	1.5	1.6	1.7	1.5	1.6	1.5	1.5	1.4	1.4
Total	15.9	11.8	10.2	9.4	9.0	8.7	7.7	7.3	6.8	6.5	6.2	5.9	5.8	5.3	5.3

Units: kilotonnes

¹ "Other" for PM₁₀ includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-44 Northern Ireland PM₁₀ Emissions by Sector, 1990-2010



Northern Ireland's PM₁₀ emissions have declined by 67% since 1990 and accounted for 5% of the UK total in 2010. 18% of PM₁₀ emissions in Northern Ireland come from road transport (1A3b) sources (down by 15% since 1990), whilst 37% stem from commercial, residential and agricultural combustion (mainly of coal and solid fuels), down by 77% since 1990. Emissions from power generation (1A1a) were 17% of the total emissions in 1990, but have been reduced to 1% of the Northern Ireland total in 2010. Per capita emissions of

PM₁₀ in Northern Ireland are 60% higher than the UK average due to the high contribution of emissions from residential combustion and agricultural mobile machinery.


Figure 2-45 Map of PM₁₀ Emissions in Northern Ireland, 2010



2.6 SULPHUR DIOXIDE

Since 1970 there has been a substantial overall reduction of more than 94% in SO₂ emissions from across the UK, mainly due to a decline in emissions from combustion of sulphur-containing solid fuels and petroleum products.

Figure 2-46 UK Sulphur Dioxide Emissions by Constituent Country, 1990-2010



Emissions from combustion of petroleum products have fallen significantly due to the decline in fuel oil use and the reduction in the sulphur content of gas oil and DERV (diesel fuel specifically used for road vehicles). The reduced sulphur content of gas oil is particularly significant in sectors such as domestic heating, commercial heating and off-road sources where gas oil is used extensively.

Fuel combustion accounts for 92% of total UK SO₂ emissions with the sulphur arising from the fuel itself. The SO₂ emission can be calculated from knowledge of the sulphur content of the fuel and from information on the amount of sulphur retained in the ash. Published fuel consumption data (DECC, 2011a), sulphur contents of liquid fuels (Tucker, 2011) and data from coal producers regarding sulphur contents of coals enable reliable estimates to be produced. The main combustion sources are:

- **Power generation (1A1a).** Power stations account for 43% of UK SO₂ emissions in 2010. Historically coal-fired stations have been the most important source, but the gradual change in fuel mix of UK power stations (to more nuclear and gas-fired plant) and improvements in generation efficiency and these reductions will continue in the near future as more CCGT stations are built and FGD is fitted to more coal-fired power plant.
- **Industrial Combustion (1A2).** Emissions of SO₂ from industry result from the combustion of coal and oil, some refinery processes and the production of sulphuric acid and other chemicals. Between 1970 and 2009 emissions from industrial combustion sources have fallen by 96%, primarily due to the decline in energy-intensive heavy industries such as iron & steel manufacturing. In addition, UK industry has gradually switched from coal and oil-based fuels in favour of natural gas, as it provides a cleaner, cheaper energy source.

In 2010, road transport (1A3b) emissions account for less than 1% of the total SO₂ emissions. Previously this source was more significant, but a tightening of fuel standards during the 1990s and more recently due to the EU Fuel Quality Directive and its amendments has led to a significant decline in emissions due to the reduction

in the sulphur content of DERV. The reduction in the sulphur content of gas oil (to 0.1% by mass from January 2008 onwards) has also reduced emissions from off-road vehicles.

Emissions from domestic (1A4b), commercial & institutional sectors (1A4a) have also declined since 1970, reflecting the major changes in fuel mix from oil and coal to gas. Emissions from waste incinerators have reduced significantly during the 1990s due to the introduction of stricter emission standards forcing the closure of old-design incinerators and their replacement with more modern plant with improved abatement.

Emissions of SO₂ for England, Wales, Scotland and Northern Ireland are summarised in the tables and graphs below. **Table 2-31** shows how total UK SO₂ emissions are split between the four constituent countries, and **Table 2-32** shows per capita emissions.

Table 2-31 Proportion of UK Sulphur Dioxide Emissions by Constituent Country

Year	England	Scotland	Wales	Northern Ireland	Unallocated
1990	84%	8%	5%	3%	0%
2010	68%	20%	8%	4%	1%

Table 2-32 Emissions of Sulphur Dioxide per capita by Constituent Country (kg/head)

Year	England	Scotland	Wales	Northern Ireland	UK
1990	64.9	58.8	65.4	71.1	64.8
2010	5.3	15.5	10.9	8.0	6.5

2.6.1 England Sulphur Dioxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the SO₂ emissions in England by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-33 England Emissions of Sulphur Dioxide by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	2480.7	1470.1	1011.6	722.9	712.2	638.1	593.6	596.8	456.8	356.3	315.5	271.9	208.6	152.5	149.9
Industrial Combustion	331.1	241.4	148.7	114.2	108.4	108.7	94.8	86.7	89.2	91.3	80.8	75.9	71.4	63.9	59.5
Transport Sources	70.4	58.8	35.8	27.8	20.2	18.0	18.4	18.7	18.6	18.9	18.5	13.9	12.3	11.1	10.5
Commercial, domestic and agricultural combustion	129.9	95.0	69.3	63.6	50.7	48.5	35.5	31.7	29.3	25.2	23.4	23.1	24.0	22.0	22.1
Other ¹	85.6	78.4	74.3	63.5	52.5	49.2	43.5	45.8	47.8	46.5	45.5	50.4	41.6	33.1	33.1
Total	3097.7	1943.6	1339.7	992.1	944.0	862.4	785.8	779.7	641.7	538.3	483.8	435.3	358.0	282.7	275.1

Units: kilotonnes


¹ "Other" for sulphur dioxide includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-47 England Sulphur Dioxide Emissions by Sector, 1990-2010



England's SO₂ emissions have declined by 91% since 1990 and accounted for 68% of the UK total in 2010. Power generation (1A1a) is by far the most significant source, accounting for 39% of the England total in 2010 (mainly from the sulphur in coal and fuel oil), but due to the growth in gas & nuclear fuel use and the installation of FGD plant at a number of coal-fired power stations, emissions from this source have reduced by 95% since 1990. 22% of SO₂ emissions in England are from industrial combustion (1A2: down by 82% since 1990), 14% from refineries (1A1b: down 61% since 1990) whilst residential combustion and national navigation contribute 6% and 3% of the total respectively. Reductions in SO₂ emissions across all sectors are also due to the progress towards production of low-sulphur petroleum-based fuels such as gas oil (diesel) and burning oil.

Figure 2-48 Map of Sulphur Dioxide Emissions in England, 2010



2.6.2 Scotland Sulphur Dioxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the SO₂ emissions in Scotland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-34 Scotland Emissions of Sulphur Dioxide by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	217.0	135.0	96.5	81.4	104.3	101.0	96.3	82.3	67.0	53.8	66.1	53.1	46.9	49.7	65.7
Industrial Combustion	37.0	21.6	9.9	7.0	6.6	6.9	9.6	9.2	9.7	10.8	8.7	7.6	5.0	3.6	3.2
Transport Sources	11.4	9.8	7.0	6.4	5.3	4.8	5.0	4.5	4.4	4.4	4.0	2.9	2.6	2.4	2.2
Commercial, domestic and agricultural combustion	28.7	24.5	20.8	19.4	16.6	15.4	13.0	11.7	10.5	9.2	8.7	8.4	9.0	8.0	8.5
Other ¹	4.7	1.7	1.7	1.9	1.7	2.7	2.0	1.8	1.8	1.7	1.4	1.7	1.6	1.4	1.4
Total	298.8	192.6	135.9	116.1	134.5	130.7	125.9	109.6	93.5	79.9	88.9	73.7	65.1	65.2	81.0

Units: kilotonnes

¹ "Other" for sulphur dioxide includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-49 Scotland Sulphur Dioxide Emissions by Sector, 1990-2010



Scotland's SO₂ emissions have declined by 73% since 1990 and account for 20% of the UK total. Power generation (1A1a) is by far the most significant source, accounting for 73% of the Scotland total in 2010 (mainly from the sulphur in coal and fuel oil), but due to the growth in gas, renewable and nuclear fuel use, emissions from this source have reduced by 70% since 1990. However, in recent years, fluctuating trends in electricity generation have had a noticeable impact on emissions of SO₂ from power generation (1A1a) in Scotland. In 2006, coal-fired generation in Scotland reached a peak, producing 17,488 GWh since then it has declined by 16% in 2010 (down to 14,653 GWh). The trends in coal-powered generation are reflected in the emissions of SO₂ from the power sector in Scotland, which showed an increase in 2006 in comparison to preceding and following years (Figure 2-49). (Note, in the table and figure above the Energy Industries sector includes power generation, petroleum refining and other energy industries such as collieries and gas processing.)

In 2010, 4% of SO₂ emissions in Scotland are from industrial combustion (1A2: down by 91% since 1990), 8% from refineries (1A1b: down 70% since 1990) whilst residential combustion and national navigation contribute 10% and 2% of the total respectively. The overall downward trend in SO₂ emissions across all sectors is also due to the progressive introduction of low-sulphur petroleum-based fuels such as gas oil (diesel) and burning oil.

Figure 2-50 Map of Sulphur Dioxide Emissions in Scotland, 2010



2.6.3 Wales Sulphur Dioxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the SO₂ emissions in Wales by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-35 Wales Emissions of Sulphur Dioxide by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	108.8	69.7	51.8	40.2	58.8	51.7	42.2	46.3	44.2	39.2	45.8	37.2	24.3	17.4	16.8
Industrial Combustion	41.9	30.3	19.0	16.0	16.1	16.2	10.3	8.8	6.2	4.6	4.2	5.2	5.0	4.6	5.3
Transport Sources	5.9	5.0	3.5	3.0	2.5	2.2	2.2	2.2	2.4	2.4	2.2	1.6	1.5	1.5	1.6
Commercial, domestic and agricultural combustion	17.4	13.7	10.5	10.0	8.0	7.6	5.6	5.1	4.7	3.9	3.7	3.9	4.1	3.8	3.8
Other ¹	13.1	11.9	12.1	11.2	8.9	7.6	4.9	5.1	5.4	5.8	6.9	6.9	6.0	3.9	5.4
Total	187.1	130.6	96.8	80.4	94.3	85.2	65.2	67.4	63.0	55.9	62.8	54.8	40.9	31.2	32.9

Units: kilotonnes

¹ "Other" for sulphur dioxide includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-51 Wales Sulphur Dioxide Emissions by Sector, 1990-2010



Wales SO₂ emissions have declined by 82% since 1990 and accounted for 8% of the UK total in 2010. In 2010, emissions from petroleum refineries are the most significant source in Wales, accounting for 37% of all SO₂ emissions (1A1b: down 46% since 1990). Previously the power generation (1A1a) sector was the highest emitter (40% of Wales emissions in 2007, and 50% in 2006) but the installation of Flue Gas Desulphurisation at the coal-fired Aberthaw station has significantly reduced the emissions from power generation to only 12% of the Wales total in 2010 (a reduction in emissions of 82% between 2007 and 2010).

In 2010, 16% of SO₂ emissions in Wales are from industrial combustion (1A2: down by 87% since 1990), 10% from residential combustion and 19% from iron and steel production (1A2a + 2C1: down by 40% since 1990). Reductions in SO₂ emissions across all sectors are also due to the progress towards production of low-sulphur petroleum-based fuels such as gas oil (diesel) and burning oil.

Figure 2-52 Map of Sulphur Dioxide Emissions in Wales, 2010



2.6.4 Northern Ireland Sulphur Dioxide Inventory by Sector, 1990-2010

The table and graph below give a summary of the SO₂ emissions in Northern Ireland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-36 Northern Ireland Emissions of Sulphur Dioxide by Sector, 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	68.2	39.7	26.8	26.8	28.3	29.9	18.3	17.4	16.4	14.9	13.5	8.3	11.0	4.6	2.3
Industrial Combustion	18.9	15.7	8.2	5.1	4.2	4.6	2.7	2.4	2.8	3.9	3.7	3.4	3.5	3.0	3.4
Transport Sources	2.5	2.3	1.5	1.3	1.0	0.9	0.9	1.0	1.0	1.0	1.0	0.7	0.7	0.6	0.6
Commercial, domestic and agricultural combustion	23.5	21.7	19.0	17.1	15.4	13.8	12.5	11.2	9.7	8.7	8.2	7.8	8.2	7.3	7.8
Other ¹	0.3	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2
Total	113.4	79.6	55.7	50.5	49.0	49.3	34.6	32.1	30.1	28.6	26.5	20.4	23.6	15.7	14.3

Units: kilotonnes


¹ "Other" for sulphur dioxide includes emissions from Fugitive, Industrial Processes, Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-53 Northern Ireland Sulphur Dioxide Emissions by Sector, 1990-2010



Northern Ireland's SO₂ emissions have declined by 87% since 1990 and they accounted for 4% of the UK total in 2010. Residential combustion is by far the most significant source accounting for 51% of the Northern Irish total in 2010 (1A4bi: down 57% since 1990) which is much higher than the rest of the UK, reflecting the higher use of coal and solid fuels in the domestic sector (1A4b) in this region. Power generation (1A1a) accounted for 60% of emissions in 1990 (mainly from the sulphur in coal and fuel oil), but due to the growth in gas use, emissions from this source have reduced by 97% since 1990. 24% of SO₂ emissions in Northern Ireland are from industrial combustion (1A2: down by 82% since 1990). These emissions are expected to decline in the future as the gas supply network develops further and solid fuel use is reduced. Reductions in SO₂ emissions across all sectors are due to the use of low-sulphur petroleum-based fuels such as gas oil (diesel) and burning oil.


Figure 2-54 Map of Sulphur Dioxide Emissions in Northern Ireland, 2010



2.7 LEAD

Since 1970, emissions of lead to air in the UK have declined by 99%, with reductions in emissions occurring in most sectors. However, the decline has been mainly driven by the progressive phasing out of leaded petrol in the UK. The lead content of leaded petrol was reduced from around 0.34 g/l to 0.143 g/l in 1986. From 1987, sales of unleaded petrol increased, particularly as a result of the increased use of cars fitted with three-way catalyst and leaded petrol was then phased out from general sale at the end of 1999. The largest source of lead until 1999 was the road transport sector (1A3b).

Figure 2-55 UK Lead Emissions by Constituent Country, 1990-2010



In recent years, the main sources of lead emissions in the UK are from iron and steel combustion, metal production and combustion of lubricants in industry. There has been some reduction in emissions from iron and steel production processes due to improved abatement measures. Emissions have also declined as a result of the decreasing use of coal.

- Iron and Steel Sector Production (2C1).** This sector is responsible for 38% of the lead emissions in the UK in 2010, of which sinter production accounts for 74%. Emissions from sinter production have reduced by around 63% since 1990, but the proportion of annual UK lead emissions it accounts for has increased due to the decline in emissions from sectors such as transport (1A3).
- Industrial Processes (2).** Emissions of lead from industry arise only from a few industries, dominated by metal production. Other sources include the chemical industry and solid fuel transformation. Between 1970 and 2010 emissions from metal production and the chemical industry have fallen by 91% and 96% respectively.
- Waste Incineration (6C).** In 1990, lead emissions from waste incineration accounted for 8% (238t) of the overall emissions in the UK. Municipal Solid Waste (MSW) incinerators not meeting regulatory standards were closed in the period leading up to December 1996. Improved combustion and flue gas controls, and developments in abatement technology in modern incinerator design has resulted in emissions from waste incineration declining down to less than 2% of the UK total.

In 2010, transport (1A3) emissions accounted for 3% of the total lead emissions. Previously this source was the most significant, accounting for 74% of emissions in 1990. Emissions of lead for England, Wales, Scotland and Northern Ireland are summarised in the tables and graphs below. **Table 2-37** shows how total UK lead emissions are split between the four constituent countries and **Table 2-38** shows per capita emissions.

Table 2-37 Proportion of UK Lead Emissions by Constituent Country

Year	England	Scotland	Wales	Northern Ireland	Unallocated
1990	86%	7%	5%	2%	0%
2010	68%	6%	23%	3%	0%

Table 2-38 Emissions of Lead per capita by Constituent Country (g/head)

Year	England	Scotland	Wales	Northern Ireland	UK
1990	51.8	41.2	47.4	44.9	50.4
2010	0.8	0.7	4.5	0.9	1.0

2.7.1 England Lead Inventory by Sector, 1990-2010

The table and graph below give a summary of the lead emissions in England by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-39 England Emissions of Lead by Sector (tonnes), 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	137.9	117.0	17.8	14.0	13.9	11.3	10.2	10.1	10.1	9.1	9.0	2.6	2.6	2.6	2.7
Industrial Combustion	31.4	25.3	19.7	16.6	13.8	14.3	13.9	13.5	14.5	12.3	9.6	10.3	9.7	12.0	10.1
Transport Sources	1800.6	878.2	481.7	253.2	2.1	2.0	2.0	2.0	2.0	2.1	2.1	1.7	1.7	1.6	1.6
Commercial, domestic and agricultural combustion	22.3	12.9	10.0	9.2	6.3	6.2	4.6	4.0	3.7	3.2	3.0	3.1	3.3	3.4	3.4
Fugitive	2.8	1.9	1.9	1.7	1.6	1.7	2.1	1.7	1.9	2.0	2.1	2.3	2.3	1.9	1.9
Industrial Processes	244.9	207.6	173.0	101.0	76.9	79.1	74.4	59.3	50.7	50.6	40.4	39.6	35.0	25.5	20.0
Other ¹	230.4	68.9	0.5	0.7	0.7	0.3	1.5	1.4	1.2	0.7	0.7	0.7	0.7	0.7	0.7
Total	2470.2	1311.9	704.6	396.5	115.3	115.0	108.7	91.9	84.1	80.0	66.8	60.3	55.3	47.7	40.4

Units: tonnes

¹ "Other" for lead includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-56 England Lead Emissions by Sector, 1990-2010



England's lead emissions have declined by 98% since 1990 and accounted for 68% of the UK total in 2010. The emissions that arise due to the production in the iron and steel industries represent the most significant source, accounting for 30% of the England total in 2010. 50% of the overall emissions are from industrial processes, including chemicals, iron and steel, aluminium and other metal production. Transport (1A3) sources used to dominate the emission of lead in England in 1990, accounting for 73%. However, as

mentioned previously, due to the phase out of leaded petrol, transport sources (1A3) now only account for 4% of lead emissions in England.

Figure 2-57 Map of Lead Emissions in England, 2010



2.7.2 Scotland Lead Inventory by Sector, 1990-2010

The table and graph below give a summary of the lead emissions in Scotland by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-40 Scotland Emissions of Lead by Sector (tonnes), 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	11.84	7.57	2.71	2.20	3.00	2.78	1.92	1.52	3.35	1.58	1.69	0.76	0.44	0.66	1.30
Industrial Combustion	3.30	2.60	1.96	1.74	1.49	1.49	1.73	1.57	2.06	1.40	1.04	1.09	0.95	0.98	1.03
Transport Sources	170.76	81.69	43.82	23.12	0.25	0.23	0.23	0.23	0.23	0.23	0.23	0.20	0.19	0.19	0.18
Commercial, domestic and agricultural combustion	4.40	2.45	1.99	1.91	1.48	1.39	1.17	1.00	0.88	0.73	0.75	0.78	0.86	0.86	0.90
Fugitive	0.47	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Industrial Processes	11.16	5.12	2.87	2.20	1.78	1.82	1.95	1.13	0.91	0.77	0.65	0.78	0.45	0.30	0.27
Other ¹	7.35	2.55	0.04	0.06	0.06	0.03	0.16	0.14	0.12	0.07	0.07	0.08	0.07	0.07	0.07
Total	209.28	101.99	53.39	31.22	8.06	7.74	7.15	5.58	7.55	4.79	4.43	3.67	2.97	3.05	3.76

Units: tonnes

¹ "Other" for lead includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-58 Scotland Lead Emissions by Sector, 1990-2010



Scotland's lead emissions have declined by 98% since 1990 and accounted for 6% of the UK total in 2010. In Scotland, emission from energy industries is the most significant source, accounting for 34% of the Scotland total in 2010 (1A1: down 89% since 1990). Emissions from power generation (1A1a) accounted for 30% but this is an increase in proportion of the overall emissions relative to 1990 when it was only responsible for 6%

of overall emissions. This is due to the decreased emissions from transport sources (1A3: down more than 99% since 1990). Emissions from domestic combustion accounted for 20% of the Scotland total emissions in 2010 (1A4b: down 80% since 1990), whilst other industrial combustion accounted for 25% (1A2f: down 66% since 1990).

Figure 2-59 Map of Lead Emissions in Scotland, 2010



2.7.3 Wales Lead Inventory by Sector, 1990-2010

The table and graph below give a summary of the lead emissions in Wales by broad NFR sector categories. The disaggregation of these categories is available in Appendix A (see Sector Category column).


Table 2-41 Wales Emissions of Lead by Sector (tonnes), 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	3.71	1.79	0.83	0.58	1.46	1.01	0.49	0.82	0.37	0.69	0.81	0.40	0.46	0.48	0.45
Industrial Combustion	3.00	2.44	1.91	2.50	2.37	1.74	4.79	3.36	7.80	2.90	0.94	1.00	0.90	0.95	1.05
Transport Sources	110.03	52.84	27.69	14.53	0.14	0.13	0.13	0.13	0.14	0.14	0.14	0.12	0.12	0.11	0.11
Commercial, domestic and agricultural combustion	2.68	1.78	1.49	1.50	1.11	1.11	0.82	0.71	0.66	0.54	0.55	0.60	0.65	0.67	0.69
Fugitive	1.45	1.82	1.81	1.82	1.99	1.39	0.51	0.80	0.43	0.41	0.44	0.29	0.21	0.21	0.26
Industrial Processes	14.39	16.39	16.77	15.81	15.79	11.37	6.27	10.94	14.54	16.17	12.16	11.22	10.85	8.63	10.91
Other ¹	0.44	0.37	0.02	0.03	0.03	0.01	0.09	0.08	0.07	0.04	0.04	0.04	0.04	0.04	0.04
Total	135.70	77.42	50.51	36.77	22.90	16.76	13.10	16.84	24.02	20.89	15.08	13.67	13.22	11.10	13.52

Units: tonnes

¹ "Other" for lead includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.


Figure 2-60 Wales Lead Emissions by Sector, 1990-2010



Wales lead emissions have declined by 90% since 1990 and accounted for 23% of the UK total in 2010. The emissions that arise due to the production in the iron and steel industries are by far the most significant source, accounting for 77% of the Wales total in 2010. 81% of lead emissions in Wales are from industrial production (2: down by 24% since 1990), 8% from industrial combustion and 2% fugitive emissions from fuels (1A2 down 65% and 1B down 81% since 1990), and 3% arise from energy industries (1A1: down by 88% since 1990) and transport sources (1A3: down more than 99% since 1990). The proportion of emissions from the

industrial combustion sector is explained by the above average concentration of heavy industry within the country. This is the main reason that reductions in lead emissions in Wales are less than those achieved in England, Scotland, and Northern Ireland, and is also the reason per capita emissions are much higher in Wales than the other constituent countries in 2010.

Figure 2-61 Map of Lead Emissions in Wales, 2010



2.7.4 Northern Ireland Lead Inventory by Sector, 1990-2010

The table and graph below give a summary of the lead emissions in Northern Ireland by broad NFR sector categories.


Table 2-42 Northern Ireland Emissions of Lead by Sector (tonnes), 1990-2010

	1990	1995	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Energy Industries	2.77	1.31	0.68	0.21	0.24	0.22	0.18	0.15	0.05	0.05	0.06	0.03	0.01	0.02	0.05
Industrial Combustion	1.62	1.42	1.13	1.02	0.84	0.85	0.92	0.75	0.76	0.80	0.63	0.67	0.59	0.61	0.68
Transport Sources	62.89	31.67	16.99	9.18	0.08	0.07	0.08	0.08	0.08	0.08	0.08	0.07	0.07	0.07	0.07
Commercial, domestic and agricultural combustion	3.47	2.06	1.67	1.51	1.31	1.13	1.05	0.88	0.70	0.59	0.56	0.56	0.62	0.60	0.65
Fugitive	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Industrial Processes	0.61	0.53	0.31	0.28	0.30	0.29	0.27	0.14	0.16	0.13	0.11	0.13	0.10	0.07	0.07
Other ¹	0.23	0.20	0.01	0.02	0.02	0.01	0.05	0.05	0.04	0.02	0.02	0.02	0.02	0.02	0.02
Total	71.60	37.19	20.79	12.22	2.79	2.58	2.54	2.04	1.80	1.67	1.46	1.48	1.42	1.39	1.54

Units: tonnes


¹ "Other" for lead includes emissions from Solvent Processes, Agriculture, Waste as well as 1A5b, 7A.

Figure 2-62 Northern Ireland Lead Emissions by Sector, 1990-2010



Northern Ireland's lead emissions have declined by 98% since 1990 and accounted for 3% of the UK total in 2010. In Northern Ireland, emissions that arise from domestic combustion (1A4b) account for 36% of the Northern Ireland total in 2010 (1A4b: down 81% since 1990). 44% of lead emissions in Northern Ireland come from industrial combustion (1A2: down 58% since 1990), 4% from transport sources (1A3: down more than 99% since 1990) and 3% arise from energy industries (1A1: down by 98% since 1990).

Figure 2-63 Map of Lead Emissions in Northern Ireland, 2010



3 Uncertainties

As discussed in Section 1.2, the air quality pollutant inventories for England, Scotland, Wales and Northern Ireland are derived using a “top-down” approach whereby the UK inventory totals are disaggregated across the four countries. For most sources there is insufficient regional data to enable a comprehensive “bottom-up” calculation to be made, and hence available proxy data are used to estimate the country-specific share of UK activity for each emission source.

The calculated uncertainties of the UK inventories for AQ pollutants are shown in the table below:

Table 3.1 Uncertainty calculated for the UK Emission Inventories of AQ Pollutants

Pollutant	Estimated Uncertainty %
PM ₁₀	- 20 to + 30
Carbon Monoxide	+/- 20
Oxides of Nitrogen	+/- 10
Sulphur Dioxide	+/- 4
Non-Methane Volatile Organic Compounds	+/- 10
Ammonia	+/- 20
Lead	-20 to +30

(Source: “UK Informative Inventory Report (1980 to 2010)”, Passant *et al.*, 2012)

Further to these uncertainties in the UK datasets, there is an additional uncertainty inherent in the methodologies of disaggregating the UK emissions across the four countries. Further to this, there is greater uncertainty for emission estimates in the early years of the time series, as these estimates are frequently based on very limited historic data; in more recent years the development of environmental regulation and reporting has increased, for example through the development of annual reporting of emissions by operators of major industrial plant, regulated under Environmental Permitting Regulations (EPR) / Integrated Pollution Prevention and Control (IPPC).

The air quality pollutant inventories for England, Scotland, Wales and Northern Ireland are therefore subject to greater uncertainty than the equivalent UK estimates. The level of uncertainty is anticipated to reduce as further research is conducted and more data reporting at local and regional level is developed; one example of this is the on-going DECC sub-national energy statistics work programme, which includes annual research tasks targeted to improve local and regional energy data in specific source sectors.

The key characteristics of each inventory are discussed below, by pollutant, with an indicative “Uncertainty Rating” provided in each case.

3.1 AMMONIA

Ammonia emission estimates are more uncertain than SO₂, NO_x and NMVOC inventories due largely to the nature of the major agricultural sources. Emissions depend on animal species, age, weight, diet, housing systems, waste management and storage techniques and environmental conditions. Hence emissions are affected by a large number of factors that make the interpretation of experimental data difficult and emission estimates uncertain (DOE, 1994). Emission estimates for non-agricultural sources such as wild animals are also highly uncertain. Unlike the case of NO_x and NMVOC, a few sources dominate the inventory and there is limited potential for error compensation.

Uncertainty Rating: HIGH

3.2 CARBON MONOXIDE

In 2010, 43% of the UK total CO came from road transport sources (1A3b) alone, with 46% of UK carbon monoxide emissions derived from other sources of fuel combustion. Emission estimates for road transport are highly uncertain, as the available dataset of emission measurements is small and shows significant variability. Emissions from stationary combustion processes are also variable and depend on the technology employed and the specific combustion conditions. Emission estimates from small and medium-sized installations are derived from emission factors based on relatively few measurements of emissions from different types of boiler. As a result of the high uncertainty in major sources, emission estimates for CO are much more uncertain than other pollutants such as NO_x, CO₂ and SO₂ which are also emitted mainly from combustion processes. Unlike the case of NO_x and NMVOC, a few sources dominate the inventory and there is limited potential for error compensation.

Uncertainty Rating: HIGH

3.3 NITROGEN OXIDES

NO_x emission estimates are less accurate than SO₂ because they are calculated using measured emission factors, which can vary widely with combustion conditions; emission factors given in the literature for combustion sources show large variations. In the case of road transport (1A3b) emissions, while the inventory methodology takes into account variations in the amount of NO_x emitted as a function of speed and vehicle type, significant variations in measured emission factors have been found even when keeping these parameters constant.

From the above, one might expect the NO_x inventory to be very uncertain, however the overall uncertainty is in fact lower than any pollutant other than SO₂ for a number of reasons:

- While NO_x emission factors may be somewhat uncertain, activity data used in the NO_x inventory is very much more certain. This contrasts with inventories for pollutants such as volatile organic compounds and PM₁₀, which contain a higher degree of uncertainty.
- The NO_x inventory is made up of a large number of emission sources with many of similar size and with none dominating. This leads to a large potential for error compensation, where an underestimate in emissions in one sector is very likely to be compensated by an overestimate in emissions in another sector.
- Many of the larger point-source emission sources make up the bulk of the regional estimates, and these are commonly derived from extrapolation of on-line measurement data and hence are regarded to be good quality.

Uncertainty Rating: LOW

3.4 NON-METHANE VOLATILE ORGANIC COMPOUNDS

The NMVOC inventory is more uncertain than SO₂ and NO_x inventories. This is due in part to the difficulty in obtaining good emission factors or emission estimates for some sectors (e.g. fugitive sources of NMVOC emissions from industrial processes, and natural sources) and partly due to the absence of good activity data for some sources. As with NO_x, there is a high potential for error compensation, and this is responsible for the relatively low level of uncertainty compared with most other pollutants in the NAEI.

Uncertainty Rating: MODERATE

3.5 PM₁₀

The UK emission inventory for PM₁₀ has undergone considerable revision over recent years through specific research into key source sectors to improve the veracity of emission factors and improve the “bottom-up” activity data such as fuel use. Nonetheless, the uncertainties in the PM₁₀ emission estimates must still be considered high, due to persisting uncertainties in some sectors regarding emission factors, activity data and particulate size distribution profiles.

Emission factors are generally based on a few measurements on an emitting source that is assumed to be representative of all similar sources. Emission estimates for PM₁₀ are based whenever possible on source-specific measurements of PM₁₀, but frequently the available data is emission measurement of total particulate matter and hence conversion to PM₁₀ is required based either on the size distribution of the sample collected or (more usually) on literature data on typical size distributions.

Many sources of particulate matter are diffuse or fugitive in nature, such as emissions from coke ovens, metal processing, raw material stockpiles, loading and unloading activities, construction or quarrying sites. These emissions are difficult to measure and are often dependent on conditions that vary over time and between localities such as meteorology and topography and hence are also difficult to model accurately. In many such cases it is likely that no satisfactory estimates or measurements have ever been made.

Emission estimates for combustion of fuels are generally considered more reliable than those for industrial processes, quarrying and construction. All parts of the inventory would need to be improved before the overall uncertainty could be reduced to the levels seen in the inventories for SO₂, NO_x, or NMVOC.

Uncertainty Rating: HIGH

3.6 SULPHUR DIOXIDE

Sulphur dioxide emissions can be estimated with most confidence as they depend largely on the level of sulphur in fuels. Hence the DA inventories, being based upon comprehensive analysis of coals and fuel oils consumed by power stations and the agriculture, industry and domestic sectors, contain accurate emission estimates for the most important sources.

Uncertainty Rating: LOW

3.7 LEAD

The lead inventory is more uncertain than SO₂ and NO_x inventories, and the certainty of the emissions varies over the time-series as different source sectors dominate at different times due to the very significant reductions in emissions from the key sources in 1990, notably road transport (1A3b). From the key sources in 1990, the lead emission estimates were based on measured concentrations of lead in the fuels, which were tightly regulated prior to being phased out in the late 1990s. This gives a high confidence in the estimates for those sources of fuel combustion, which dominated in the early 1990s, but are now much reduced.

In more recent years, the level of emissions is estimated to be very much lower, and derived from a smaller number of sources. The metal processing industries are mainly regulated under IPPC and the estimates provided by plant operators to the regulatory agencies and used in the national inventories are based on emission measurements or emission factors that have been researched for the specific process type. There is a moderate level of uncertainty associated with these annual emission estimates due to the discrete nature of the stack emissions monitoring techniques and determination of mass emission flow rates from point sources. Furthermore the variability of lead content of raw materials such as fuels (e.g. coal) is such that the discrete lead emission measurements provide a snap-shot of the process and plant performance, and there is some uncertainty as regards how representative that result may be for use in scaling up to provide annual emission

estimates. These uncertainties are inherent within the inventories from environmental regulators of IPC/IPPC industries and are unavoidable; the emissions data from IPPC regulated installations used in the compilation of these DA inventories are subject to a managed process of quality checking by the environmental regulatory agencies and are regarded as the best data available for inventory compilation.

The observed year-to-year variations in emission estimates are based on actual trends reported by plant operators and may reflect changes in lead content of raw materials. The uncertainty in emission monitoring applies to all pollutants to some degree, but more so for pollutants such as lead for which (i) no continuous emission monitoring systems are available, and (ii) where fuel composition is known to be highly variable depending on the fuel source. This is not the case for species such as NO_x and SO₂ where many regulated sites will use Continuous Emission Monitoring Systems and the fuel elemental composition is either not a significant factor in process emissions or does not vary as much as for heavy metals and other trace contaminants. The emission estimates of lead from other smaller-scale combustion and process sources from industrial and commercial activities are less well documented and the estimates are based on emission factors that are less certain than those based on regulatory emissions monitoring and reporting.

Uncertainty Rating: MODERATE

4 References

- Agricultural Industries Confederation (2006). Fertiliser Statistics 2006 Report www.agindustries.org.uk
- Alcan (2004, 2011), Personal Communication.
- APEG (1999), Source Apportionment of Airborne Particulate Matter in the United Kingdom. Report of the Airborne Particles Expert Group. ISBN 0-7058-1771-7
- ATOC (2011), Fuel use by train operating companies, 2011, personal communication.
- AQEG (2005). "Particulate Matter in the United Kingdom", *Air Quality Expert Group*, <http://archive.defra.gov.uk/environment/quality/air/airquality/publications/particulate-matter/documents/pm-summary.pdf>
- Bell, DM (2011), ISR and Power Station Fuel Consumption, Personal Communication, Northern Ireland Department of Environment.
- BGS (2011), United Kingdom Minerals Yearbook 1990-2010. British Geological Survey.
- Boulter, P.G., Latham, S. (2009). "Emission Factors 2009: Report 4 – A Review of Methodologies for Estimating Cold Start Emissions", *TRL Project Report PPR 357*, June 2009. <http://www.dft.gov.uk/pgr/roads/environment/emissions/report-4.pdf>
- Boulter PG, TJ Barlow and IS McRae (2009). "Emission Factors 2009: Report 3 - Exhaust Emission Factors for Road Vehicles in the UK", *TRL Project Report PPR 356*, June 2009. <http://www.dft.gov.uk/pgr/roads/environment/emissions/report-3.pdf>
- British Cement Association (2011), Cement industry fuel use data and plant capacity data, Personal Communication.
- British Glass (2011), Production Statistics for UK Glass Industry, Personal Communication.
- CAA (2011). "UK airport statistics 2010" – annual, CAA.
- Carslaw, D., Beevers, S., Westmoreland, E., Williams, M., Tate, J., Murrells, T., Stedman, J., Li, Yvonne, Grice, S., Kent. A., and Tsagatakis, I (2011). "Trends in NO_x and NO₂ emissions and ambient measurements in the UK", *Report for Defra* http://uk-air.defra.gov.uk/reports/cat05/1108251149_110718_AQ0724_Final_report.pdf
- Coal Authority (2011), Regional Coal Production: open cast and deep mined, Personal Communication.
- DECC (2010a). "Digest of UK Energy Statistics", *Department of Energy & Climate Change*, HMSO
- DECC (2011a), Local gas use data by LDZ, *personal communication*
- DECC (2011b). "Energy Trends December 2011", Articles on DA electricity generation and consumption patterns and the regional energy statistics for 2005 to 2010
- DECC (2011c). Oil & Gas flaring and venting volume data by installation and gas landings information, *personal communication*
- DECC Offshore Inspectorate (2011), Environmental Database for annual Emissions and Discharges from Offshore Installations, from the EEMS reporting inventory.

DECC Oil & Gas (2011), spreadsheet of emissions data from the EEMS reporting system of site operator annual returns to DECC for 2010. Personal Communication.

Defra (2001). "The Air Quality Strategy for England, Scotland, Wales and Northern Ireland" - A Consultation Document on Proposals for Air Quality Objectives for Particles, Benzene, Carbon Monoxide and PAHs, *HMSO*

Defra (2005), "The Air Quality Strategy for England, Scotland, Wales and Northern Ireland: Addendum". *HMSO*

Defra (2007), "The Air Quality Strategy for England, Scotland, Wales and Northern Ireland (Volume 1)", *HMSO*

Defra (2011), "Emissions of Air Pollutants in the UK, 1970 to 2010", *Department for Environment, Food and Rural Affairs*, December 2011. <http://www.defra.gov.uk/statistics/files/Statistical-Release-Emissions-of-air-pollutants-in-the-UK-1970-2010-UK.pdf>

Defra (2012). Revised DA waste landfill model outputs, using available DA-specific data on NSW and C&I waste. Steve Nelson, Defra, 2012.

DETR (1998a). "Review of the United Kingdom National Air Quality Strategy" - A Consultation Document. 98EP0541/A. *The Stationary Office, Norwich*

DETR (2000). "The Air Quality Strategy for England, Scotland, Wales and Northern Ireland- Working Together for Clean Air", *The Stationary Office, Norwich*

DFPNI (2010) Public Sector Energy Campaign data, personal communication

DfT (2009). Personal communication with Chris Parkin, Cleaner Fuels and Vehicles, Transport Statistics, DfT, March 2009

DfT (2011a) Vehicle Licensing Statistics: 2010, Transport Statistics Bulletin SB(10)16, 2011.

DfT (2011b) "Transport Statistics Great Britain: 2010 edition", The Stationary Office

DfT (2011c) "Road Statistics 2010: Traffic, Speeds and Congestion" Transport Statistics Bulletin (SB (10) 18), 2011

DfT (2011d) Transport Statistics Great Britain, Table 2.2. Air Transport Movements

DfT (2011e), Road Freight Statistics 2010, Transport Statistics Bulletin SB(10) 20, August 2011.

DOE (1994). "Impacts of Nitrogen Deposition on Terrestrial Ecosystems", Report of The UK Review Group on Impacts of Atmospheric Nitrogen, ISBN 1-870393-22-8

DOE (1997). "The United Kingdom National Air Quality Strategy", March 1997, (Cm 3587)

DRDNI (2011a) "Northern Ireland Transport Statistics 2009-2010", Central Statistics and Research Branch, Department of Regional Development in Northern Ireland. http://www.drdni.gov.uk/ni_transport_statistics_2009-10.pdf

DRDNI (2011b), "Traffic and Travel Information Report, September 2011, Incorporating Annual Traffic Census and Vehicle Kilometres of Travel"

DRDNI (2011c), personal communication with Leah Ginn, Central Statistics and Research Branch, Department for Regional Development Northern Ireland, September 2011.

EEA (2010a), EMEP/CORINAIR Emission Inventory Guidebook – 2009. Technical report No 9/2009. <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009>

EEA (2010b), EMEP/EEA Emission Inventory Guidebook – 2009. Part B: sectoral guidance chapters, 1.A.3.b Road transport updated June 2010. European Environment Agency.

<http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009>

Environment Agency (2011a), database of emissions to atmosphere from authorised processes in England & Wales, as reported to the Pollution Inventory (data for 1998-2010).

Environment Agency (2011b), database of EUETS operator data including fuel use, process details and emissions totals, personal communication.

EPAQS (1995), Expert Panel on Air Quality Standards – Particles. HMSO. London.

Fernando, S. (2011). Update of Estimated Methane Emissions from UK Abandoned Coal Mines. At http://uk-air.defra.gov.uk/reports/cat07/1107080945_1775-ghg-improvement-project-wsp-report.pdf

Firmus Energy (2011). Gas sales in Northern Ireland in 2010: domestic and non-domestic data, Personal Communication

ISSB (2011). Iron & Steel Industry Annual Statistics for the UK, including regional fuel use data. Personal Communication, Donna Leach

MacCarthy J, Thomas J, Choudrie S, Pearson B, S. Passant, N. Thistlethwaite, G. Murrells, T. Watterson J. Cardenas, Malcolm H (2011). UK Greenhouse Gas Inventory, 1990 to 2009. AEA ISBN 0-9554823-9-9

Misselbrook, T.H., Sutton, M.A., Scholefield, D., (2004). "A simple process-based model for estimating ammonia emissions from agricultural land after fertilizer applications", *Soil Use and Management* 20, 365-372.

T P Murrells, N R Passant, G Thistlethwaite, A Wagner, Y Li, T Bush, J Norris, C Walker, R A Stewart, I Tsagatakis, R Whiting, C Conolly, S Okamura, M Peirce, S Sneddon, J Webb, J Thomas, J MacCarthy, C Martinez, S Choudrie, N Brophy (2011a), UK Emissions of Air Pollutants 1970-2009. AEA, Harwell, Oxfordshire.

National Grid (2011), natural gas leakage from high pressure, low pressure distribution systems and from Above Ground Installations, personal communication

NIEA (2011a), spreadsheet of emissions to atmosphere from authorised processes in Northern Ireland, as reported to the Inventory of Statutory Releases (data for 2005-2010), *Northern Ireland Environment Agency*

NIEA (2011b), spreadsheet of EU ETS operator data including fuel use, process details and emissions totals, personal communication, *Northern Ireland Environment Agency*

Northern Gas Networks (2011). Natural Gas leakage from LDZ and AGIs, personal communication

ONS (2011). Datasets available pertaining to a wide range of industrial and population-related activities via the publications "Annual Abstract of Statistics" and "Regional Trends", including regional GDP data, *Office for National Statistics*

Pang, Y., Walker, H., and Murrells, T (2011). "Improving the Greenhouse Gas Inventories for Road Transport in Scotland, Wales and Northern Ireland", Report for DECC, Welsh Assembly Government, the Scottish Government and the Department of the Environment for Northern Ireland, AEAT/ENV/R/3167, April 2011. http://uk-air.defra.gov.uk/reports/cat07/1106161316_IP_Task_23_DA_Road_Transport_Issue_1.pdf

N R Passant, T P Murrells, G Thistlethwaite, A Wagner, Y Li, T Bush, J Norris, R Whiting, C Walker, R A Stewart, I Tsagatakis, C Conolly, N C J Brophy, S Okamura (2012). "UK Informative Inventory Report (1980 to 2010)"
Phoenix Natural Gas (2007), Gas consumption detailed by end-user sectors for 2005 and gas leakage data, Personal Communication

Phoenix Natural Gas (2011), Gas consumption and gas leakage data for 2010 and breakdown of gas sales by customer sectors 2006-2010, Personal Communication

QUARG (1996), Airborne Particulate Matter in the United Kingdom, Third report of the Quality of Urban Air Review Group. Department of the Environment. ISBN 0-9520771-3-2.

Rothamsted Research (2011). Ammonia emissions from agricultural sources and livestock data by region, personal communication (Tom Misselbrook).

Scotia Gas Networks (2011) Natural Gas leakage from LDZ and AGIs, personal communication

SEPA (2011a), Annual atmospheric emissions data for authorised processes in Scotland, from SEPA's Pollution Release Inventory, Personal Communication.

SEPA (2011b), spreadsheet of EU ETS operator data including fuel use, process details and emissions totals, personal communication.

Tata Steel (2011). Site-specific breakdown of pollutant emissions by sub-source, for all UK integrated steelworks, *personal communication* (Peter Quinn)

Thistlethwaite, G., Goodwin, J., Salisbury, E., MacCarthy, J., Pang, Y., Thomson, A., Cardenas, L., (2012). "Greenhouse Gas Inventories for England, Scotland, Wales and Northern Ireland: 1990 – 2010", http://uk-air.defra.gov.uk/reports/cat07/1207171006_DA_GHGI_report_2010_Issue1.pdf

Thomson, A.M., Hallsworth, S., Malcolm, H. (2012). Emissions and Removals of GHGs from Land Use Land Use Change and Forestry for England, Scotland, Wales and Northern Ireland: 1990-2010. At http://uk-air.defra.gov.uk/reports/cat07/1204120924_DA_LULUCF_GHG_inventory_report_2012_fullreport_v2.pdf

Tomlinson (2010) Peat use within Scotland and Northern Ireland: summary of research findings from CEH, personal communication.

Transco (2011) Historic Gas Demands by Load Category, Personal Communication

Translink (2011), Annual fuel consumption, Personal Communication.

Tsagatakis *et al.* (2011). "NAEI UK Emission Mapping Methodology 2009", AEA

Tucker (2011). "Sulphur content of liquid fuels and petrol properties", *Personal Communication*

UKPIA (2011), Pollutant emissions for UK refineries, from combustion and process sources. Personal Communication, United Kingdom Petroleum Industry Association Ltd

Vayu Ltd. (2011), Northern Ireland gas sales for domestic and commercial & industrial customers for 2010, Personal Communication.

Wales & West Utilities (2011) Natural Gas leakage from LDZ and AGIs, personal communication

Watson M (2011), Personal communication regarding sulphur content of liquid fuels, UKPIA.

Webb, J., Misselbrook, T.H., (2004). "A mass-flow model of ammonia emissions from UK livestock production", *Atmospheric Environment*, 38, 2163-2176

www.wastedataflow.org, website providing regional waste disposal fate data, to inform regional estimates of landfill waste activities.